Contents

| Appendix A | Public Scoping and Review Comments and DOE Responses | A .1 |
|------------|--|-------------|
| Pt 1 | Public Scoping Comments and Responses for the ILAW SEIS | A.1 |
| Pt 2 | Public Scoping Comments and Responses for the HSW EIS | A.20 |
| A.1 | DOE Programmatic/Nationwide Analysis | A.21 |
| | A.1.1 Coordination with Other Federal Reports, Environmental Impact, and | |
| | DOE Policy Statements | A.21 |
| | A.1.2 Nationwide Impact Comparisons and Equity Issues | A.23 |
| A.2 | Alternatives and Activities Analyzed in the HSW EIS | A.24 |
| | A.2.1 Alternative Options | A.25 |
| | A.2.1.2 Use of Commercial or Offsite Disposal Facilities | A.26 |
| | A.2.1.3 Alternative Actions and Emerging Technologies | A.27 |
| | A.2.2 Recommended Alternative Analyses | A.29 |
| A.3 | Waste Types and Volumes | A.30 |
| A.4 | Environmental Consequences and Analysis Methods | A.32 |
| A.5 | Public Involvement and Government Agency Consultations | A.34 |
| A.6 | References | A.35 |
| | | |
| Appendix B | Detailed Alternative Descriptions, Assumptions, Waste Volumes, and Waste | |
| | Stream Flowsheets | B.1 |
| B.1 | Introduction | B.1 |
| B.2 | Waste Stream Numbers | B.1 |
| B.3 | HSW EIS Waste Processing Assumptions | |
| B.4 | Waste Volumes | B.9 |
| B.5 | Radionuclide Inventories. | B.15 |
| B.6 | Waste Stream Flowsheets | B.49 |
| B.7 | References | B.102 |
| Appendix C | Description of Waste Volumes for the Hanford Site Solid (Radioactive and | |
| ** | Hazardous) Waste Program EIS | C.1 |
| C.1 | Volume Identification, Review, and Selection Methodology | |
| C.2 | Low-Level Waste | |
| C.3 | Mixed Low-Level Waste | |
| C.4 | Transuranic Waste | C.9 |
| C.5 | Waste Treatment Plant Wastes | C.11 |
| C.6 | References | C.12 |
| Appendix D | Supplemental Information on the Low Level Burial Grounds, Environmental | |
| ** | Restoration Disposal Facility, Borrow Pits, Trench Liners, and Disposal | |
| | Facility Barriers | D.1 |
| D.1 | Low Level Burial Grounds | D.1 |
| | D.1.1 200 East Area Burial Grounds | D.1 |
| | D.1.2 200 West Area Burial Grounds | D.4 |
| D.2 | Environmental Restoration Disposal Facility | D.11 |
| D.3 | Borrow Pit Resource Excavation | D.12 |
| D 4 | Liner Ontions for Disposal Facilities | D 16 |

| D.5 | Barrie | Options. | | D.17 |
|------------|--------|------------|--|------|
| | D.5.1 | Hanford | Barrier | D.18 |
| | D.5.2 | | l RCRA Subtitle C Barrier | |
| | D.5.3 | | d RCRA Subtitle D Barrier | |
| | D.5.4 | | ual Cover Barrier with Bentonite Mix | |
| D.6 | | • | | |
| Appendix E | Air Qu | ality Anal | lysis | E.1 |
| E.1 | | | ine Emissions | |
| E.2 | | _ | | |
| E.3 | | | imum Air Quality Impacts | |
| E.4 | | | eneral Conformity Review | |
| E.5 | | | | E.26 |
| Appendix F | Metho | ds for Eva | luating Impacts on Health from Radionuclides and Chemicals | F.1 |
| F.1 | | | on Impact Assessment Methods | |
| | F.1.1 | • | t Releases to the Atmosphere | |
| | 1.1.1 | F.1.1.1 | Release Fractions for the Waste Receiving and | 1.2 |
| | | 1 .1.1.1 | Processing Facility | F.15 |
| | | F.1.1.2 | Release Fractions for the Existing T Plant Complex | |
| | | F.1.1.3 | The New Waste Processing Facility and Modified T Plant | 1.17 |
| | | 1.1.1.3 | • | F.21 |
| | | T: 1 1 4 | Complex | |
| | E 1 0 | F.1.1.4 | Pulse Drier Operation | |
| | F.1.2 | | Point Characteristics | |
| | F.1.3 | • | heric Transport | |
| | F.1.4 | • | e Scenarios | |
| | | F.1.4.1 | Industrial Scenario | |
| | | F.1.4.2 | Resident Gardener Scenario | F.34 |
| | | F.1.4.3 | Soil (Air or Irrigation Water Deposition) | |
| | | | Transport Medium | |
| | | F.1.4.4 | Air Transport Medium | F.38 |
| | | F.1.4.5 | Waterborne Transport Medium | F.38 |
| | F.1.5 | Soil Acc | cumulation Model | F.39 |
| | | F.1.5.1 | Evaluation of Distribution Coefficient for | |
| | | | Organic Chemicals | F.41 |
| | | F.1.5.2 | Evaluation of Distribution Coefficients for Radionuclides | |
| | | | and Inorganic Chemicals | F.42 |
| | F.1.6 | Health I | mpacts | |
| | F.1.7 | | r Radiological Health Consequences | |
| | F.1.8 | | ison of Radiation Risk Results for Children—Estimated | |
| | 1.1.0 | | ederal Guidance Reports 11 and 13 | F.45 |
| F.2 | Accide | • | Assessment Methods | F.47 |
| 1.2 | F.2.1 | | nent Method | |
| | F.2.1 | | t Frequency | |
| | | | A • | |
| | F.2.3 | | diological Impact Endpoints | |
| F 2 | т. • | F.2.3.1 | Impacts from Industrial Accidents | |
| F.3 | | | Assessment Methods | |
| | F.3.1 | Human | Intrusion Exposure Scenarios | F.52 |

| | F.3.2 | Radiolog | gical Analysis | F.54 |
|------------|---------|--------------------|---|--------------|
| F.4 | | | terborne Pathways | F.55 |
| F.5 | | | Impacts of West Valley TRU Wastes Processed or Stored | |
| | | | | F.18 |
| | F.5.1 | | Health Impacts | F.18 |
| | F.5.2 | | Atmospheric Release – Public Health Impacts | F.18 |
| F.6 | | | | F.18 |
| 1.0 | 1101010 | 11000 | | 1.10 |
| Appendix G | Ground | dwater Ou | ality Impacts | G.1 |
| G.1 | | | l Approach | G.3 |
| 0.1 | G.1.1 | | Analysis | G.3 |
| | G.1.2 | | Analytical Approach | G.4 |
| | G.1.3 | | Ferm Release | G.7 |
| | 0.1.5 | G.1.3.1 | Constituent Grouping and Screening | G.8 |
| | | G.1.3.1 G.1.3.2 | Source Inventories | G.12 |
| | | G.1.3.2 G.1.3.3 | Release Models | G.12 |
| | G.1.4 | | Zone Modeling | G.20 |
| | 0.1.4 | G.1.4.1 | Stratigraphy | G.20 |
| | | G.1.4.1 G.1.4.2 | Hydraulic Properties | G.21 |
| | | G.1.4.2 G.1.4.3 | • | G.24 |
| | | G.1.4.3 G.1.4.4 | Recharge Rates | |
| | | G.1.4.4 G.1.4.5 | | G.24 |
| | C 1.5 | | Vadose Zone Model Implementation | G.25 G.26 |
| | G.1.5 | | vater Modeling | |
| | | G.1.5.1 | Conceptual Model | G.26 |
| | | G.1.5.2 | Simulation of Post-Closure Flow Conditions | G.36 |
| α | D () | G.1.5.3 | Simulation of Unit Releases | G.39 |
| G.2 | | | water Quality Impact Results | G.58 |
| | G.2.1 | | ve Group A | G.58 |
| | | | Wastes Disposed of Before 1996 | G.59 |
| | | | Wastes Disposed of After 1995 | G.60 |
| | G.2.2 | | ve Group B | G.61 |
| | | G.2.2.1 | Wastes Disposed of Before 1996 | G.62 |
| | | G.2.2.2 | Wastes Disposed of After 1995 | G.62 |
| | G.2.3 | | ve Group C | G.63 |
| | | G.2.3.1 | Wastes Disposed of Before 1996 | G.64 |
| | | G.2.3.2 | Wastes Disposed of After 1995 | G.64 |
| | G.2.4 | Alternati | ve Group D ₁ | G.64 |
| | | G.2.4.1 | Wastes Disposed of Before 1996 | G.65 |
| | | G.2.4.2 | Wastes Disposed of After 1995 | G.65 |
| | G.2.5 | Alternati | ve Group D ₂ | G.66 |
| | | G.2.5.1 | Wastes Disposed of Before 1996 | G.67 |
| | | G.2.5.2 | Wastes Disposed of After 1995 | G.67 |
| | G.2.6 | Alternati | ve Group D ₃ | G.68 |
| | | G.2.6.1 | Wastes Disposed of Before 1996 | G.69 |
| | | G.2.6.2 | Wastes Disposed of After 1995 | G.69 |
| | G.2.7 | Alternati | ve Group E ₁ | G.70 |
| | | G.2.7.1 | Wastes Disposed of Before 1996 | G.71 |
| | | | Wastes Disposed of After 1995 | G.71 |

| | G.2.8 | Aiternati | ve Group E_2 | G./3 |
|------------|-----------------------|-----------|--|------------|
| | | G.2.8.1 | Wastes Disposed of Before 1996 | G.73 |
| | | G.2.8.2 | Wastes Disposed of After 1995 | G.73 |
| | G.2.9 | | ve Group E ₃ | |
| | | | Wastes Disposed of Before 1996 | |
| | | | Wastes Disposed of After 1995 | |
| | G 2 10 | | on Alternative | |
| | 0.2.10 | | Wastes Disposed of Before 1996 | |
| | | | Wastes Disposed of After 1995 | |
| G.3 | Lica of i | | formance Assessment Calculations in Potential HSW EIS | 0.70 |
| 0.5 | | | ndwater Quality and Human Health Impacts | G.266 |
| | • | | | 0.200 |
| | G.3.1 | - | Waste Form and Engineering Performance Evaluated in the | 0.26 |
| | $\alpha \circ \alpha$ | | AW Performance Assessment | |
| | G.3.2 | | al Planned Analyses of Waste Disposal System Performance | |
| | G.3.3 | | Scaling of ILAW PA Results for Use in the Analysis | |
| | | G.3.3.1 | Scaling for Estimated Inventory | |
| | | G.3.3.2 | Scaling for Alternative HSW EIS Disposal Site Locations | |
| G.4 | | | ng Assumptions on Long-Term Cover System Performance | G.272 |
| G.5 | | | water Quality Impacts at Low-Level Waste Management Area | |
| | | | elected Alternatives | |
| | G.5.1 | | ale Lines of Analysis | |
| | G.5.2 | | erm Release and Vadose Zone Transport | |
| | G.5.3 | Unit-Rele | ease Calculations and Transport in Groundwater | G.277 |
| ` | G.5.4 | Summary | y of Results | G.283 |
| | | G.5.4.1 | Alternative Group D ₁ | G.293 |
| | | G.5.4.2 | Alternative Group D ₂ | G.296 |
| | | G.5.4.3 | Alternative Group D ₃ | G.297 |
| | | G.5.4.4 | Summary of Ratios to Benchmark MCLs for | |
| | | | Technetium-99 and Iodine-129 | G.297 |
| G.6 | Potentia | al Ground | water Quality Impacts from Hazardous Chemicals in | |
| | | | | G.301 |
| | G.6.1 | | y Estimates | |
| | G.6.2 | | nant Group and Screening Analysis | |
| | G.6.3 | | Methods and Other Key Assumptions | |
| | G.6.4 | | y of Results | |
| G.7 | | • | , 01100010 | |
| G.7 | Referen | 1005 | | 0.517 |
| Appendix H | Traffic | and Trans | portation | H.1 |
| Н.1 | | | ethods | H.2 |
| 11.1 | Н.1.1 | | ical Impact Analysis Methodology | |
| | H.1.2 | | (Non-Radiological) Incident-Free Risks | H.6 |
| | H.1.3 | • | iological Accident Risks in Transit | |
| | H.1.4 | | | |
| H.2 | | | us Chemical Impact Analysis | H.7 H.8 |
| п.2 | | | ping Data | |
| | H.2.1 | | nipping Data | H.9 |
| | H.2.2 | | hipping Data | H.18 |
| | H.2.3 | Accident | Risk Input Data | H.23 |

| H.3 | Results | s of Transp | portation Impact Analysis | H.29 | | | |
|------------|---------------------------------------|-------------|--|--------------|--|--|--|
| | H.3.1 | Results of | of Onsite Transportation Impact Analysis | H.30 | | | |
| | | H.3.1.1 | Alternative Group A | H.30 | | | |
| | | H.3.1.2 | Alternative Group B | H.30 | | | |
| | | H.3.1.3 | Alternative Group C | H.30 | | | |
| | | H.3.1.4 | Alternative Group D. | H.31 | | | |
| | | H.3.1.5 | Alternative Group E | H.31 | | | |
| | | H.3.1.6 | No Action Alternative | H.31 | | | |
| | | H.3.1.7 | Summary of Transportation Impacts for the Hanford Only | | | | |
| | | | Waste Volume | H.38 | | | |
| | H.3.2 | Results of | of Offsite Transportation Impact Analysis | H.40 | | | |
| | | H.3.2.1 | Potential Population Radiological Impacts of | | | | |
| | | | Offsite Shipments | H.41 | | | |
| | | H.3.2.2 | Potential Non-Radiological Impacts of Offsite Shipments | H.44 | | | |
| | | H.3.2.3 | Results of the Maximally Exposed Individual Impact Analysis. | H.47 | | | |
| | H.3.3 | | y of Potential Impacts of Onsite and Offsite Waste Shipments | H.49 | | | |
| | 11.5.5 | H.3.3.1 | Hanford Solid Waste Management Lifecycle | 11.17 | | | |
| | | 11.3.3.1 | Transportation Impacts | H.49 | | | |
| | | H.3.3.2 | Sensitivity Studies | H.56 | | | |
| H.4 | Impact | | porting Construction and Capping Materials | H.62 | | | |
| H.5 | | | ic | H.66 | | | |
| H.6 | | | npacts of Offsite Shipments Within Washington and Oregon | H.67 | | | |
| 11.0 | H.6.1 | | gical Incident-Free Exposure and Accident Impact | 11.07 | | | |
| | 11.0.1 | - | Parameters | H.67 | | | |
| | H.6.2 | | liological Impact Analysis Parameters | H.70 | | | |
| | H.6.3 | | Results | H.70 | | | |
| H.7 | | | dous Chemical Impact Analysis | H.73 | | | |
| H.8 | | | s of Sabotage or Terrorist Attack | H.76 | | | |
| H.9 | | | ISW EIS Transportation Impacts to Those in | 11.70 | | | |
| 11.7 | Other Environmental Impact Statements | | | | | | |
| | H.9.1 | | son to the Final Waste Management Programmatic | 11.70 | | | |
| | 11.7.1 | | nental Impact Statement for Managing Treatment, Storage, | | | | |
| | | | posal of Radioactive and Hazardous Waste | H.79 | | | |
| | H.9.2 | _ | son to the Waste Isolation Pilot Plant Disposal Phase | 11.77 | | | |
| | 11.7.2 | | pplemental Environmental Impact Statement | H 81 | | | |
| H.10 | Effects | | porting Solid Waste by Rail | H.84 | | | |
| H.11 | Refere | • | orting bond waste by Kun | | | | |
| 11.11 | IXCICIO. | nees | | 11.00 | | | |
| Appendix I | Ecolog | rical Resor | irces | I.1 | | | |
| I.1 | | | inces | I.1 | | | |
| I.2 | | | strial Resources Resulting from Surface Disturbance | I.2 | | | |
| 1.2 | I.2.1 | | ve Group A | I.2 | | | |
| | I.2.2 | | ve Group B | I.13 | | | |
| | I.2.2 I.2.3 | Alternati | ve Group C | I.15 | | | |
| | I.2.3 I.2.4 | | ve Groups D ₁ , D ₂ , and D ₃ | I.16 | | | |
| | I.2.5 | | ve Groups E_1 , E_2 , and E_3 | I.10 I.17 | | | |
| | I.2.5 I.2.6 | | on Alternative | I.17 I.18 | | | |
| | I.2.6 I.2.7 | | on Alternative | I.18 I.19 | | | |
| | 1.4./ | wingane | Л1 | 1.19 | | | |

| | I.2.8 Biodiversity | I.27 |
|------------|--|--------------|
| | I.2.9 Microbiotic Crusts | I.28 |
| I.3 | Potential Impacts to Columbia River Aquatic and Riparian Resources Resulting | |
| | from Future Contaminant Releases | I.28 |
| | I.3.1 Assumptions Regarding Contaminants | I.29 |
| | I.3.2 Assumptions Regarding Partitioning of Contaminants to Abiotic Media | I.30 |
| | I.3.3 Ecological Contaminant Exposure Model | I.31 |
| | I.3.4 Combined Radiological Toxicity | I.31 |
| | I.3.5 Chemical Toxicity of Total Uranium | I.45 |
| I.4 | Consultations | I.55 |
| I.5 | References | I.55 |
| 1.5 | References | 1.55 |
| Appendix J | Construction Noise – Method of Assessment | J.1 |
| J.1 | Assessment of Noise Impacts | J.1 |
| | J.1.1 Critical Distances | J.1 |
| | J.1.2 Source Term | J.2 |
| J.2 | References | J.3 |
| 3.2 | TCTCTCTCCCS | 3.5 |
| Appendix K | Cultural Resources | K.1 |
| K.1 | Introduction | K.1 |
| | K.1.1 200 East and 200 West Areas | K.1 |
| | K.1.2 Central Waste Complex Expansion Area | K.5 |
| | K.1.3 New Waste Processing Facility | |
| K.2 | Area C – Borrow Pits, Stockpile Area, and Access Roads | |
| | K.2.1 Literature and Record Search – Previous Cultural Resources Surveys | K.7 |
| | K.2.2 Research Initiatives and Field Reconnaissance | K.8 |
| K.3 | References | K.50 |
| | | |
| Appendix L | System Assessment Capability: A 10,000-Year Post-Closure Assessment | L.1 |
| L.1 | Introduction | L.1 |
| | L.1.1 Context of SAC Runs | L.3 |
| | L.1.2 Relationship to EIS Calculations | L.4 |
| L.2 | Methods and Approach | L.5 |
| | L.2.1 Modular Components of SAC | L.6 |
| | L.2.2 Inventory | L.8 |
| | L.2.2.1 Initial Assessment Inventory | L.10 |
| | L.2.2.2 Comparison of HSW EIS and Initial Assessment Inventories | L.10 |
| | L.2.3 Release | L.15 |
| | L.2.3.1 Conceptual Model | L.16 |
| | L.2.3.2 Implementation Model | L.16 |
| | L.2.3.3 Numerical Models Relevant to HSW EIS | L.17 |
| | L.2.3.4 Comparison of Release Model Parameters | L.21 |
| | L.2.4 Vadose Zone Module | L.23 |
| | L.2.4.1 Distribution Coefficients (K _d s) for Technetium-99, | L.43 |
| | Iodine-129, and Uranium | L.23 |
| | L.2.4.2 Vadose Zone Strata and Hydraulic Properties | L.25 |
| | L.2.4.2 Vadose Zone Strata and Hydraunic Properties | L.23 L.27 |
| | L.2.4.3 Surface Covers | L.27 |
| | | |

| | L.2.6 River Transport Module | L.34 |
|------------|--|--------------|
| | L.2.7 Risk and Impact | L.35 |
| | L.2.8 Uncertainty | L.36 |
| L.3 | Results | L.36 |
| | L.3.1 Release to Groundwater Results | L.37 |
| | L.3.2 Drinking Water Dose at Selected 200 East and 200 West Area Locations. L.3.2.1 Drinking Water Dose at the Northeast Corner of the | L.74 |
| | 200 West Area | L.75 |
| | L.3.2.2 Drinking Water Dose at the Southeast Corner of the | |
| | 200 East Area | L.76 |
| | L.3.2.3 Drinking Water Dose at the Northwest Corner of the 200 East Area | L.79 |
| | L.3.3 Dose from Columbia River Water at the City of Richland Pumping Station | |
| | L.3.3.1 Drinking Water Dose at the City of Richland Pumping Station. | L.82 |
| | L.3.4 Annual Drinking Water Dose at Selected 200 East Area and Columbia | L.84 |
| L.4 | River Locations from Hanford Sources Including ILAW | L.02 L.94 |
| L.4 | References | L.92 |
| Appendix M | Long-Term Impacts Associated with Discontinuing Disposal of HSW at the | |
| | Hanford Site | M.1 |
| M.1 | Introduction | M.1 |
| M.2 | Impacts on Groundwater | M.1 |
| Appendix N | Overview of DOE Nationwide and Hanford Site Waste Management Programs | |
| 11 | and Initiatives | N.1 |
| N.1 | | N.1 |
| | | N.1 |
| | · · · · · · · · · · · · · · · · · · · | N.3 |
| N.2 | | N.5 |
| | N.2.1 Spent Nuclear Fuel | N.5 |
| | N.2.2 High-Level Waste | N.5 |
| | N.2.3 Environmental Restoration Waste | N.6 |
| | N.2.4 Groundwater Protection | N.8 |
| | N.2.5 Liquid Waste | N.9 |
| | N.2.6 Cleanup, Constraints, and Challenges Team (C3T) | N.10 |
| | N.2.7 Hanford Performance Management Plan (HPMP) | N.11 |
| | - The state of the | N.15 |
| | N2.9 Transuranic Waste Considerations | N.18 |
| N.3 | References | N.19 |
| | | |
| Appendix O | Unpublished Sources Cited in the Hanford Solid (Radioactive and Hazardous) | |
| | Waste Environmental Impact Statement | Ω 1 |

Figures

| B.1 | Waste Types and Waste Streams Considered in the HSW EIS | B.2 |
|------|---|------|
| D.1 | 218-E-12B Burial Ground | D.2 |
| D.2 | 218-E-10 Burial Ground | D.3 |
| D.3 | 218-W-3A Burial Ground | |
| D.4 | 218-W-3AE Burial Ground | |
| D.5 | 218-W-4B Burial Ground | |
| D.6 | 218-W-4C Burial Ground | |
| D.7 | 218-W-5 Burial Ground | D.9 |
| D.8 | 218-W-6 Burial Ground | D.10 |
| D.9 | Existing and Proposed ERDF Disposal Cells | D.11 |
| D.10 | Area C Location Relative to the 200 East and 200 West Burial Grounds | D.12 |
| D.11 | Borrow Pit Layout in Area C | D.13 |
| D.12 | Hanford Barrier | D.19 |
| D.13 | Standard RCRA Subtitle C Barrier | |
| D.14 | Modified RCRA Subtitle D Barrier with Bentonite Mix | |
| D.15 | US Ecology, Inc. Conceptual Cover Barrier | D.22 |
| F.1 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well 1 km Downgradient from the 200 West Area, Alternative Group A | F.57 |
| F.2 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well 1 km Downgradient Northwest of 200 East Area, Alternative | |
| | Group A | F.58 |
| F.3 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well 1 km Downgradient Southeast of 200 East Area, Alternative | |
| | Group A | F.59 |
| F.4 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well Adjacent to the Columbia River Alternative Group A | F.60 |
| F.5 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well 1 km Downgradient from 200 West Area, Alternative Group B | F.61 |
| F.6 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well 1 km Downgradient Northwest of 200 East Area, Alternative | |
| | Group B | F.62 |
| F.7 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well Adjacent to the Columbia River, Alternative Group B | F.63 |
| F.8 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well 1 km Downgradient from 200 West Area, Alternative Group C | F.64 |
| F.9 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well 1 km Downgradient Northwest of 200 East Area, Alternative | |
| | Group C | F.65 |
| F.10 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well 1 km Downgradient Southeast of 200 East Area, Alternative | |
| | Group C | F.66 |
| F.11 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well Adjacent to the Columbia River, Alternative Group C | F.67 |

| F.12 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
|------|--|-------|
| | Water from a Well 1 km Downgradient from 200 West Area, Alternative Group D ₁ | F.68 |
| F.13 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well 1 km Downgradient Northwest of 200 East Area, Alternative | |
| | Group D ₁ | F.69 |
| F.14 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well 1 km Downgradient Southeast of 200 East Area, Alternative | |
| | Group D_1 | F.70 |
| F.15 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well Adjacent to the Columbia River, Alternative Group D ₁ | F.71 |
| F.16 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well 1 km Downgradient from 200 West Area, Alternative Group D ₂ | F.72 |
| F.17 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well 1 km Downgradient Northwest of 200 East Area, Alternative | |
| | Group D_2 | F.73 |
| F.18 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well Adjacent to the Columbia River, Alternative Group D ₂ | F.74 |
| F.19 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well 1 km Downgradient from 200 West Area, Alternative Group D ₃ | F.75 |
| F.20 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | 21,70 |
| 1.20 | Water from a Well 1 km Downgradient from ERDF, Alternative Group D ₃ | F.76 |
| F.21 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | 2170 |
| | Water from a Well 1 km Downgradient Northwest of 200 East Area, Alternative | |
| | Group D ₃ | F.77 |
| F.22 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well Adjacent to the Columbia River, Alternative Group D ₃ | F.78 |
| F.23 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well 1 km Downgradient from 200 West Area, Alternative Group E ₁ | F.79 |
| F.24 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well 1 km Downgradient from ERDF, Alternative Group E ₁ | F.80 |
| F.25 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well 1 km Downgradient Northwest of 200 East Area, Alternative | |
| | Group E ₁ | F.81 |
| F.26 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well Adjacent to the Columbia River, Alternative Group E ₁ | F.82 |
| F.27 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well 1 km Downgradient from 200 West Area, Alternative Group E ₂ | F.83 |
| F.28 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well 1 km Downgradient from ERDF, Alternative Group E ₂ | F.84 |
| F.29 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well 1 km Downgradient Northwest of 200 East Area, Alternative | |
| | Group E ₂ | F.85 |
| F.30 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well 1 km Downgradient Southeast of 200 East Area, Alternative | |
| | Group E ₂ | F.86 |
| F.31 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| | Water from a Well Adjacent to the Columbia River, Alternative Group E ₂ | F.87 |
| | | |

| F.32 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in Water from a Well 1 km Downgradient from 200 West Area, Alternative Group E ₃ | F.88 |
|-------|--|------|
| F.33 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | Г.00 |
| 1.33 | Water from a Well 1 km Downgradient from ERDF, Alternative Group E ₃ | F.89 |
| F.34 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | 1.07 |
| 1.51 | Water from a Well 1 km Downgradient Northwest of 200 East Area, Alternative | |
| | Group E ₃ | F.90 |
| F.35 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | 1.70 |
| 1.00 | Water from a Well 1 km Downgradient Southeast of 200 East Area, Alternative | |
| | Group E ₃ | F.91 |
| F.36 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | 1.71 |
| 1.00 | Water from a Well Adjacent to the Columbia River, Alternative Group E ₃ | F.92 |
| F.37 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | |
| 1.0, | Water from a Well 1 km Downgradient from 200 West Area, No Action Alternative | F.93 |
| F.38 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | 1.,0 |
| | Water from a Well 1 km Downgradient Northwest of 200 East Area, | |
| | No Action Alternative | F.93 |
| F.39 | Hypothetical Annual Drinking Water Dose at Various Times over 10,000 Years in | , - |
| | Water from a Well Adjacent to the Columbia River, No Action Alternative | F.94 |
| G.1 | Lines of Analysis Downgradient of Aggregate Hanford Solid Waste Disposal Areas | G.3 |
| G.2 | Schematic Representation of Computational Framework and Codes Used in this | |
| | SW EIS | G.7 |
| G.3 | Changes in Infiltration Rates Assumed in Source-Term Release to Approximate | |
| | the Modified RCRA Subtitle C Barrier System Degradation | G.17 |
| G.4 | STOMP Code Results for Releases to the Water Table for a Unit Release from | |
| | LLBGs for an Assumed Recharge Rate of 0.5 cm/yr | G.26 |
| G.5 | Comparison of Generalized Hydrogeologic and Geologic Stratigraphy | G.28 |
| G.6 | Peripheral Boundaries Defined for the Three-Dimensional Model | G.30 |
| G.7 | Transmissivity Distribution for the Unconfined Aquifer System Based on | |
| | Two Dimensional Inverse Model Calibration | G.32 |
| G.8 | Distribution of Estimated Hydraulic Conductivities at Water Table from Best-Fit | |
| | Inverse Calibration of Sitewide Groundwater Model | G.34 |
| G.9 | Distribution of Estimated Hydraulic Conductivities Along Section Lines A-A' | |
| | and B-B' from Best-Fit Inverse Calibration of Sitewide Groundwater Model | G.35 |
| G.10 | Predicted Post-Hanford Water Table Conditions (Predominant Northerly Flow) | G.37 |
| G.11 | Predicted Post-Hanford Water Table Conditions (Predominant Easterly Flow) | G.38 |
| G.12a | Simulated Transport of a 10-year Unit Release of a Contaminant Representative of | |
| | Mobility Class 1 from MLLW in the 200 West Area at 100 Years After Release | |
| | Using a Groundwater Model with a Predominant Northerly Flow from the | |
| | Central Plateau | G.40 |
| G.12b | Simulated Transport of a 10-year Unit Release of a Contaminant Representative of | |
| | Mobility Class 1 from MLLW in the 200 West Area at 300 Years After Release | |
| | Using a Groundwater Model with a Predominant Northerly Flow from the | |
| | Central Plateau | G41 |

| G.12c | | |
|-------|---|------|
| | Mobility Class 1 from MLLW in the 200 West Area at 500 Years After Release | |
| | Using a Groundwater Model with a Predominant Northerly Flow from the | |
| | Central Plateau | G.42 |
| G.12d | Simulated Transport of a 10-year Unit Release of a Contaminant Representative of | |
| | Mobility Class 1 from MLLW in the 200 West Area at 700 Years After Release | |
| | Using a Groundwater Model with a Predominant Northerly Flow from the | |
| | Central Plateau | G.43 |
| G.13a | Simulated Transport of a 10-Year Unit Release of a Contaminant Representative of | |
| 0.10 | Mobility Class 1 from MLLW in the 200 East Area at 50 Years After Release Using | |
| | a Groundwater Model with a Predominant Northerly Flow from the Central Plateau | G.44 |
| G.13b | Simulated Transport of a 10-Year Unit Release of a Contaminant Representative of | 0.11 |
| 0.130 | Mobility Class 1 from MLLW in the 200 East Area at 150 Years After Release Using | |
| | | G.45 |
| C 12. | a Groundwater Model with a Predominant Northerly Flow from the Central Plateau | G.43 |
| G.13c | Simulated Transport of a 10-Year Unit Release of a Contaminant Representative of | |
| | Mobility Class 1 from MLLW in the 200 East Area at 250 Years After Release Using | C 16 |
| C 14 | a Groundwater Model with a Predominant Northerly Flow from the Central Plateau | G.46 |
| G.14a | Simulated Transport of a 10-Year Unit Release of a Contaminant Representative of | |
| | Mobility Class 1 from MLLW in the 200 West Area at 100 Years After Release Using | ~ 4= |
| ~ | a Groundwater Model with a Predominant Easterly Flow from the Central Plateau | G.47 |
| G.14b | Simulated Transport of a 10-Year Unit Release of a Contaminant Representative of | |
| | Mobility Class 1 from MLLW in the 200 West Area at 300 Years After Release Using | |
| | a Groundwater Model with a Predominant Easterly Flow from the Central Plateau | G.48 |
| G.14c | Simulated Transport of a 10-Year Unit Release of a Contaminant Representative of | |
| | Mobility Class 1 from MLLW in the 200 West Area at 500 Years After Release Using | |
| | a Groundwater Model with a Predominant Easterly Flow from the Central Plateau | G.49 |
| G.14d | Simulated Transport of a 10-Year Unit Release of a Contaminant Representative of | |
| | Mobility Class 1 from MLLW in the 200 West Area at 700 Years After Release Using | |
| | a Groundwater Model with a Predominant Easterly Flow from the Central Plateau | G.50 |
| G.15a | Simulated Transport of a 10-Year Unit Release of a Contaminant Representative of | |
| | Mobility Class 1 from MLLW in the 200 East Area at 50 Years After Release Using | |
| | a Groundwater Model with a Predominant Easterly Flow from the Central Plateau | G.51 |
| G.15b | Simulated Transport of a 10-Year Unit Release of a Contaminant Representative of | |
| | Mobility Class 1 from MLLW in the 200 East Area at 150 Years After Release Using | |
| | a Groundwater Model with a Predominant Easterly Flow from the Central Plateau | G.52 |
| G.15c | • | |
| 0.100 | Mobility Class 1 from MLLW in the 200 East Area at 250 Years After Release Using | |
| | a Groundwater Model with a Predominant Easterly Flow from the Central Plateau | G.53 |
| G.16a | Comparison of Predicted Concentrations from Unit Releases from the 200 East Area at | 0.55 |
| G.10a | the 200 East LOAs Using Groundwater Models with a Predominant Northerly and | |
| | Easterly Flow from the Central Plateau | G.54 |
| G.16b | Comparison of Predicted Concentrations from Unit Releases from the 200 East Area at | 0.54 |
| 0.100 | | |
| | Columbia River LOAs Using Groundwater Models with a Predominant Northerly and | C 55 |
| C 17a | Easterly Flow from the Central Plateau Comparison of Prodicted Concentrations from Unit Polesces from the 200 West Area at | G.55 |
| G.17a | Comparison of Predicted Concentrations from Unit Releases from the 200 West Area at | |
| | the 200 West and ERDF LOAs Using Groundwater Models with a Predominant | 0.51 |
| | Northerly and Easterly Flow from the Central Plateau | G.56 |

| Comparison of Predicted Concentrations from Unit Releases from the 200 West Area at | |
|---|--|
| the Columbia River LOAs Using Groundwater Models with a Predominant Northerly | |
| and Easterly Flow from the Central Plateau | G.57 |
| Tc-99 and I-129 Concentration Profiles at the 1-km Lines of Analysis | |
| · | G.189 |
| · · · · · · · · · · · · · · · · · · · | |
| · | G.190 |
| · · · · · · · · · · · · · · · · · · · | 0.170 |
| · · · · · · · · · · · · · · · · · · · | G 191 |
| · · · · · · · · · · · · · · · · · · · | 0.171 |
| $oldsymbol{\cdot}$ | G 102 |
| · · · · · · · · · · · · · · · · · · · | 0.192 |
| | C 102 |
| <u>*</u> | G.193 |
| · · · · · · · · · · · · · · · · · · · | C 104 |
| | G.194 |
| | C 105 |
| | G.195 |
| | G 10. |
| | G.196 |
| | ~ 40= |
| | G.197 |
| | ~ |
| | G.198 |
| | |
| | G.199 |
| | |
| | G.200 |
| | |
| | G.201 |
| U-238 and C-14 Concentration Profiles at the 1-km Lines of Analysis (200 East) | |
| (Alternative Group A – Upper Bound Volume Wastes Disposed of After 1995) | G.202 |
| U-238 and C-14 Concentration Profiles at the 1-km Line of Analysis (200 West) | |
| (Alternative Group A – Upper Bound Volume Wastes Disposed of After 1995) | G.203 |
| Tc-99 and I-129 Concentration Profiles at the 1-km Lines of Analysis (200 East) | |
| (Alternative Group B – Hanford Only Wastes Disposed of After 1995) | G.204 |
| Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (200 West) | |
| (Alternative Group B – Hanford Only Wastes Disposed of After 1995) | G.205 |
| I-129 and Tc-99 Concentration and River Flux Profiles Near the Columbia River | |
| (Alternative Group B – Hanford Only Wastes Disposed of After 1995) | G.206 |
| Tc-99 and I-129 Concentration Profiles at the 1-km Lines of Analysis (200 East) | |
| (Alternative Group B – Upper Bound Volume Wastes Disposed of After 1995) | G.207 |
| Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (200 West) | |
| (Alternative Group B – Upper Bound Volume Wastes Disposed of After 1995) | G.208 |
| I-129 and Tc-99 Concentration and River Flux Profiles Near the Columbia River | |
| | G.209 |
| | |
| (Alternative Group C – Hanford Only Wastes Disposed of After 1995) | G.210 |
| | the Columbia River LOAs Using Groundwater Models with a Predominant Northerly and Easterly Flow from the Central Plateau |

| G.40 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (200 West) | |
|------|---|-------|
| | (Alternative Group C – Hanford Only Wastes Disposed of After 1995) | G.211 |
| G.41 | I-129 and Tc-99 Concentration and River Flux Profiles Near the Columbia River | |
| | (Alternative Group C – Hanford Only Wastes Disposed of After 1995) | G.212 |
| G.42 | Tc-99 and I-129 Concentration Profiles at the 1-km Lines of Analysis (200 East) | |
| | (Alternative Group C – Upper Bound Volume Wastes Disposed of After 1995) | G.213 |
| G.43 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (200 West) | |
| | (Alternative Group C – Upper Bound Volume Wastes Disposed of After 1995) | G.214 |
| G.44 | I-129 and Tc-99 Concentration and River Flux Profiles Near the Columbia River | |
| | (Alternative Group C – Upper Bound Volume Wastes Disposed of After 1995) | G.215 |
| G.45 | Tc-99 and I-129 Concentration Profiles at the 1-km Lines of Analysis (200 East) | |
| | (Alternative Group D ₁ – Hanford Only Wastes Disposed of After 1995) | G.216 |
| G.46 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (200 West) | |
| | (Alternative Group D ₁ – Hanford Only Wastes Disposed of After 1995) | G.217 |
| G.47 | I-129 and Tc-99 Concentration and River Flux Profiles Near the Columbia River | |
| | (Alternative Group D ₁ – Hanford Only Wastes Disposed of After 1995) | G.218 |
| G.48 | Tc-99 and I-129 Concentration Profiles at the 1-km Lines of Analysis (200 East) | |
| | (Alternative Group D ₁ – Upper Bound Volume Wastes Disposed of After 1995) | G.219 |
| G.49 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (200 West) | |
| | (Alternative Group D ₁ – Upper Bound Volume Wastes Disposed of After 1995) | G.220 |
| G.50 | I-129 and Tc-99 Concentration and River Flux Profiles Near the Columbia River | |
| | (Alternative Group D ₁ – Upper Bound Volume Wastes Disposed of After 1995) | G.221 |
| G.51 | Tc-99 and I-129 Concentration Profiles at the 1-km Lines of Analysis (200 East) | |
| | (Alternative Group D ₂ – Hanford Only Wastes Disposed of After 1995) | G.222 |
| G.52 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (200 West) | |
| | (Alternative Group D ₂ – Hanford Only Wastes Disposed of After 1995) | G.223 |
| G.53 | I-129 and Tc-99 Concentration and River Flux Profiles Near the Columbia River | |
| | (Alternative Group D ₂ – Hanford Only Wastes Disposed of After 1995) | G.224 |
| G.54 | Tc-99 and I-129 Concentration Profiles at the 1-km Lines of Analysis (200 East) | |
| | (Alternative Group D ₂ – Upper Bound Volume Wastes Disposed of After 1995) | G.225 |
| G.55 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (200 West) | |
| | (Alternative Group D ₂ – Upper Bound Volume Wastes Disposed of After 1995) | G.226 |
| G.56 | I-129 and Tc-99 Concentration and River Flux Profiles Near the Columbia River | |
| | (Alternative Group D ₂ – Upper Bound Volume Wastes Disposed of After 1995) | G.227 |
| G.57 | Tc-99 and I-129 Concentration Profiles at the 1-km Lines of Analysis (200 East) | |
| | (Alternative Group D ₃ – Hanford Only Wastes Disposed of After 1995) | G.228 |
| G.58 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (200 West) | |
| | (Alternative Group D ₃ – Hanford Only Wastes Disposed of After 1995) | G.229 |
| G.59 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (ERDF) | |
| | (Alternative Group D ₃ – Hanford Only Wastes Disposed of After 1995) | G.230 |
| G.60 | I-129 and Tc-99 Concentration and River Flux Profiles Near the Columbia River | |
| | (Alternative Group D ₃ – Hanford Only Wastes Disposed of After 1995) | G.231 |
| G.61 | Tc-99 and I-129 Concentration Profiles at the 1-km Lines of Analysis (200 East) | |
| | (Alternative Group D ₃ – Upper Bound Volume Wastes Disposed of After 1995) | G.232 |
| G.62 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (200 West) | |
| | (Alternative Group D ₃ – Upper Bound Volume Wastes Disposed of After 1995) | G.233 |
| G.63 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (ERDF) | |
| | (Alternative Group D ₃ – Upper Bound Volume Wastes Disposed of After 1995) | G.234 |

| G.64 | 1-129 and Tc-99 Concentration and River Flux Profiles Near the Columbia River | |
|------|---|-------|
| | (Alternative Group D ₃ – Upper Bound Volume Wastes Disposed of After 1995) | G.235 |
| G.65 | Tc-99 and I-129 Concentration Profiles at the 1-km Lines of Analysis (200 East) | |
| | (Alternative Group E ₁ – Hanford Only Wastes Disposed of After 1995) | G.236 |
| G.66 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (200 West) | |
| | (Alternative Group E ₁ – Hanford Only Wastes Disposed of After 1995) | G.237 |
| G.67 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (ERDF) | |
| | (Alternative Group E ₁ – Hanford Only Wastes Disposed of After 1995) | G.238 |
| G.68 | I-129 and Tc-99 Concentration and River Flux Profiles Near the Columbia River | |
| | (Alternative Group E ₁ – Hanford Only Wastes Disposed of After 1995) | G.239 |
| G.69 | Tc-99 and I-129 Concentration Profiles at 1-km Lines of Analysis (200 East) | |
| | (Alternative Group E ₁ – Upper Bound Volume Wastes Disposed of After 1995) | G.240 |
| G.70 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (200 West) | |
| | (Alternative Group E ₁ – Upper Bound Volume Wastes Disposed of After 1995) | G.241 |
| G.71 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (ERDF) | |
| | (Alternative Group E ₁ – Upper Bound Volume Wastes Disposed of After 1995) | G.242 |
| G.72 | I-129 and Tc-99 Concentration and River Flux Profiles Near the Columbia River | |
| | (Alternative Group E ₁ – Upper Bound Volume Wastes Disposed of After 1995) | G.243 |
| G.73 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (200 East SE) | |
| | (Alternative Group E ₂ – Hanford Only Wastes Disposed of After 1995) | G.244 |
| G.74 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (200 West) | |
| | (Alternative Group E ₂ – Hanford Only Wastes Disposed of After 1995) | G.245 |
| G.75 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (ERDF) | |
| | (Alternative Group E ₂ – Hanford Only Wastes Disposed of After 1995) | G.246 |
| G.76 | I-129 and Tc-99 Concentration and River Flux Profiles Near the Columbia River | |
| | (Alternative Group E ₂ – Hanford Only Wastes Disposed of After 1995 | G.247 |
| G.77 | Tc-99 and I-129 Concentration Profiles at the 1-km Lines of Analysis (200 East) | |
| | (Alternative Group E ₂ – Upper Bound Volume Wastes Disposed of After 1995) | G.248 |
| G.78 | Tc-99 and I-129 Concentration Profiles at 1-km Line of Analysis (200 West) | |
| | (Alternative Group E ₂ – Upper Bound Volume Wastes Disposed of After 1995) | G.249 |
| G.79 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (ERDF) | |
| | (Alternative Group E ₂ – Upper Bound Volume Wastes Disposed of After 1995) | G.250 |
| G.80 | I-129 and Tc-99 Concentration and River Flux Profiles Near the Columbia River | |
| | (Alternative Group E ₂ – Upper Bound Volume Wastes Disposed of After 1995) | G.251 |
| G.81 | Tc-99 and I-129 Concentration Profiles at the 1-km Lines of Analysis (200 East) | |
| | (Alternative Group E ₃ – Hanford Only Wastes Disposed of After 1995) | G.252 |
| G.82 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (200 West) | |
| | (Alternative Group E ₃ – Hanford Only Wastes Disposed of After 1995) | G.253 |
| G.83 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (ERDF) | |
| | (Alternative Group E ₃ – Hanford Only Wastes Disposed of After 1995) | G.254 |
| G.84 | I-129 and Tc-99 Concentration and River Flux Profiles Near the Columbia River | |
| | (Alternative Group E ₃ – Hanford Only Wastes Disposed of After 1995) | G.255 |
| G.85 | Tc-99 and I-129 Concentration Profiles at the 1-km Lines of Analysis (200 East) | |
| | (Alternative Group E ₃ – Upper Bound Volume Wastes Disposed of After 1995) | G.256 |
| G.86 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (200 West) | |
| | (Alternative Group E ₃ – Upper Bound Volume Wastes Disposed of After 1995) | G.257 |
| G.87 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (ERDF) | |
| | (Alternative Group E ₃ – Upper Bound Volume Wastes Disposed of After 1995) | G.258 |

| G.88 | 1-129 and Tc-99 Concentration and River Flux Profiles Near the Columbia River | |
|----------|---|------------------|
| | (Alternative Group E ₃ – Upper Bound Volume Wastes Disposed of After 1995) | G.259 |
| G.89 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (200 East) | |
| | (No Action Alternative - Previously Disposed of Wastes) | G.260 |
| G.90 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (200 West) | |
| | (No Action Alternative – Previously Disposed of Wastes) | G.261 |
| G.91 | I-129 and Tc-99 Concentration and River Flux Profiles Near the Columbia River | |
| | (No Action Alternative – Previously Disposed of Wastes) | G.262 |
| G.92 | Tc-99 and I-129 Concentration Profiles at the 1-km Lines of Analysis (200 East) | 0.202 |
| 0.,_ | (No Action Alternative – Hanford Only Wastes Disposed of After 1995) | G.263 |
| G.93 | Tc-99 and I-129 Concentration Profiles at the 1-km Line of Analysis (200 West) | 0.200 |
| 0.70 | (No Action Alternative – Hanford Only Wastes Disposed of After 1995) | G.264 |
| G.94 | I-129 and Tc-99 Concentration and River Flux Profiles Near the Columbia River | 0.20 |
| | (No Action Alternative – Hanford Only Wastes Disposed of After 1995) | G.265 |
| G.95 | Drinking Water Dose at a Well 100 Meters Downgradient from the ILAW | 0.200 |
| 0.70 | Disposal Facility as a Function of Time for Various ILAW Waste Form | |
| | Performance and Disposal Facility Parameters | G.267 |
| G.96 | Scaled Concentrations of Key Constituents that were Used from the | |
| | ILAW PA at the 200 East Area SE and Columbia River LOAs for Those | |
| | Alternative Groups where ILAW Disposal was Sited near the PUREX Plant, | |
| | Alternative Groups A, C, D ₁ , and E ₃ | G.270 |
| G.97 | Comparison of Predicted Peak Concentrations of Technetium-99 and | |
| | Iodine-129 at 200 East SE LOA from Upper Bound Inventories in Ungrouted | |
| | MLLW Disposed of After 2007 | G.273 |
| G.98 | Comparison of Predicted Peak Concentrations of Uranium-238 at the | |
| | 200 East SE LOA from Upper Bound Inventories in Ungrouted and Grouted | |
| | MLLW Disposed of After 2007 | G.274 |
| G.99 | Local-Scale Lines of Analysis 100 Meters Downgradient from the LLW Management | |
| | Areas in the 200 East Area | G.276 |
| G.100 | | |
| | Areas in the 200 West Area and at ERDF | G.277 |
| G.101 | Local-Scale Finite Element Grid Used in the Unit-Release Calculations in | |
| | Groundwater Beneath the 200 East Area | G.278 |
| G.102 | Local-Scale Finite Element Grid Used in Unit-Release Calculations in | |
| | Groundwater Beneath the 200 West Area | G.279 |
| G.103 | Local-Scale Finite Element Grid Used in Unit-Release Calculations in | |
| | Groundwater Beneath ERDF | G.280 |
| G.104 | Approximate Disposal Area Footprint Used in the 200 East Area to Represent | |
| | Waste Disposed of Before 2008 in the Unit-Release Calculation in Groundwater | G.281 |
| G.105 | Approximate Disposal Area Footprint Used in the 200 West Areas to Represent | |
| | Waste Disposed of Before 2008 in the Unit-Release Calculation in Groundwater | G.281 |
| G.106 | Approximate Disposal Area Footprint Used in Alternative Group D ₁ | |
| | (Near the PUREX Plant) to Represent Waste Disposed of After 2007 | |
| - | in the Unit-Release Calculation in Groundwater | G.282 |
| G.107 | Approximate Disposal Area Footprint Used in Alternative Group D ₂ | |
| | (218-E-12B LLBG) to Represent Waste Disposed of After 2007 in the | ~ ~ ~ ~ ~ |
| | Unit-Release Calculation in Groundwater | G.282 |

| | G.108 | Approximate Disposal Area Footprint used in Alternative Group D ₃ (at ERDF) to | |
|---|-------|--|-------|
| l | | Represent Waste Disposed of After 2007 in the Unit-Release Calculation in | |
| | | Groundwater | G.283 |
| | H.1 | Routes from Offsite to Hanford and from Hanford to WIPP | цээ |
| | H.2 | Illustration of Shipment Miles for Onsite and Offsite Waste Shipments | H.54 |
| ı | H.3 | Illustration of Potential Transportation Impacts of Onsite and Offsite Waste | 11.54 |
| | 11.5 | Shipments—LCFs from Radiological Incident-Free Transport, Radiological | |
| | | Accidents, and Non-Radiological Emissions | II 55 |
| l | TT 4 | | H.55 |
| ı | H.4 | Illustration of Potential Transportation Impacts of Onsite and Offsite Waste | 1156 |
| l | 11.5 | | H.56 |
| | H.5 | Transporation Routes in Washington and Oregon | H.68 |
| | I.1 | Plant Communities in Area C Before the 24 Command Fire of June 2000 | I.6 |
| | I.2 | Potential Plant Communities in Area C | I.7 |
| | I.3 | Element Occurrences of Plant Community Types in Area C | I.8 |
| | I.4 | Plant Communities in Area C After the 24 Command Fire of June 2000 | I.8 |
| | I.5 | Element Occurrences of Bitterbrush/Indian Ricegrass Sand Dune Complex Community | |
| | | Type Outside Area C in the 600 Area of Hanford Site, ALE (area west and south of | |
| | | Area C), and the Wahluke Slope (area north of the Columbia River) | I.22 |
| | I.6 | Element Occurrences of Big Sagebrush/Needle-and-Thread Grass Community | |
| | 1.0 | Type Outside Area C in the 600 Area of Hanford Site, ALE (area west and south of | |
| | | Area C), and the Wahluke Slope (area north of the Columbia River) | I.23 |
| | I.7 | Element Occurrences (on Gable Mountain and at Vernita Quarry) of Big Sagebrush/ | 1.20 |
| | 1., | Bluebunch Wheatgrass Community Type Outside Area C in the 600 Area of | |
| | | Hanford Site, ALE (area west and south of Area C), and the Wahluke Slope (area north | |
| | | of the Columbia River) | I.24 |
| | I.8 | Mayfly Radiological EHQs for Each Alternative Group in the 0- to 2,500-Year Time | 1,27 |
| | 1.0 | Period for Background Compared to the Hanford and Hanford Plus Background | |
| | | Scenarios | I.44 |
| | I.9 | Woodhouse's Toad Tadpole Radiological EHQs for Each Alternative Group in the 2,500- | 1.77 |
| | 1.7 | to 10,000-Year Time Period for Background Compared to the Hanford and Hanford Plus | |
| | | Background Scenarios | I.45 |
| | I.10 | American Coot Low and High Uranium Chemical Toxicity EHQs for Each Alternative | 1.43 |
| | 1.10 | Group in the 2,500- to 10,000-Year Time Period for Background and the Hanford | |
| | | Scenario | I.47 |
| | I.11 | American Coot Low and High Uranium Chemical Toxicity EHQs for Each Alternative | 1.4/ |
| | 1.11 | Group in the 2,500- to 10,000-Year Time Period for the Hanford Plus Background | |
| | | Scenario | I.48 |
| | I.12 | Woodhouse's Toad Tadpole Low and High Uranium Chemical Toxicity EHQs for | 1.40 |
| | 1.12 | Each Alternative Group in the 2,500- to 10,000-Year Time Period for Background | |
| | | | T 5 1 |
| | T 12 | and the Hanford Scenario. | I.51 |
| | I.13 | Woodhouse's Toad Tadpole Low and High Uranium Chemical Toxicity EHQs for Each | |
| | | Alternative Group in the 2,500- to 10,000-Year Time Period for the Hanford | 1.50 |
| | T 1 4 | Plus Background Scenario | I.52 |
| | I.14 | Carp Low and High Uranium Chemical Toxicity EHQs for Each Alternative Group in the 2.500- to 10.000-Year Time Period for Background and the Hanford Scenario | I 53 |
| | | In the 7 July to 10 thurs year time Period for Background and the Hantord Scenario | 1 7 4 |

| 1.15 | in the 2,500- to 10,000-Year Time Period for the Hanford Plus Background Scenario | I.54 |
|------|---|------|
| K.1 | Area C – Historical Features | K.7 |
| K.2 | Area C Predictive Model | K.9 |
| L.1 | Conceptual Model of the System Assessment Capability | L.8 |
| L.2 | Recharge Through Covers as a Function of Time | L.31 |
| L.3 | SAC Results for Annual Vadose Zone Release of Technetium-99 from All Solid | |
| | Waste Disposal Facilities Sites in the 200 East | L.39 |
| L.4 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Technetium-99 | |
| | from All Solid Waste Disposal Facilities Sites in the 200 East Area | L.39 |
| L.5 | SAC Results for Annual Vadose Zone Release of Technetium-99 from All Solid | |
| | Waste Disposal Facilities Sites in the 200 West Area | L.40 |
| L.6 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Technetium-99 | |
| | from All Solid Waste Disposal Facilities Sites in the 200 West Area | L.40 |
| L.7 | SAC Results for Annual Vadose Zone Release of Uranium from All Solid Waste | |
| | Disposal Facilities Sites in the 200 East | L.41 |
| L.8 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Uranium from | |
| | All Solid Waste Disposal Facilities Sites in the 200 East Area | L.41 |
| L.9 | SAC Results for Annual Vadose Zone Release of Uranium from All Solid Waste | |
| | Disposal Facilities Sites in the 200 West Area | L.42 |
| L.10 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Uranium from | |
| | All Solid Waste Disposal Facilities Sites in the 200 West Area | L.42 |
| L.11 | SAC Results for Annual Vadose Zone Release of Iodine-129 from All Solid Waste | |
| | Disposal Facility Sites in the 200 East Area | L.43 |
| L.12 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Iodine-129 from | |
| | All Solid Waste Disposal Facility Sites in the 200 East Area | L.43 |
| L.13 | SAC Results for Annual Vadose Zone Release of Iodine-129 from All Solid Waste | |
| | Disposal Facility Sites in the 200 West Area | L.44 |
| L.14 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Iodine-129 from | |
| | All Solid Waste Disposal Facility Sites in the 200 West Area | L.44 |
| L.15 | SAC Results for Annual Vadose Zone Release of Technetium-99 from All Tank | |
| | Sites in the 200 East Area | L.45 |
| L.16 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Technetium-99 | |
| | from All Tank Sites in the 200 East Area | L.45 |
| L.17 | SAC Results for Annual Vadose Zone Release of Technetium-99 from All Tank | |
| | Sites in the 200 West Area | L.46 |
| L.18 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Technetium-99 | |
| | from All Tank Sites in the 200 West Area | L.46 |
| L.19 | SAC Results for Annual Vadose Zone Release of Uranium from All Tank Sites | |
| | in the 200 East Area | L.47 |
| L.20 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Uranium from | |
| | All Tank Sites in the 200 East Area | L.47 |
| L.21 | SAC Results for Annual Vadose Zone Release of Uranium from All Tank Sites in the | |
| | 200 West Area | L.48 |
| L.22 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Uranium from | |
| | All Tank Sites in the 200 West Area | L.48 |

| L.23 | SAC Results for Annual Vadose Zone Release of Iodine-129 from All Tank Sites in | |
|--------------|--|----------|
| | the 200 East Area | L.49 |
| L.24 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Iodine-129 from | |
| | All Tank Sites in the 200 East Area | L.49 |
| L.25 | SAC Results for Annual Vadose Zone Release of Iodine-129 from All Tank Sites in | |
| | the 200 West Area | L.50 |
| L.26 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Iodine-129 from | |
| | All Tank Sites in the 200 West Area | L.50 |
| L.27 | SAC Results for Annual Vadose Zone Release of Technetium-99 from All Liquid | |
| | Discharge and Unplanned Release Sites in the 200 East Area | L.51 |
| L.28 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Technetium-99 from | |
| | All Liquid Discharge and Unplanned Release Sites in the 200 East Area | L.51 |
| L.29 | SAC Results for Annual Vadose Zone Release of Technetium-99 from All Liquid | |
| | Discharge and Unplanned Release Sites in the 200 West Area Plus SALDS | L.52 |
| L.30 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Technetium-99 from | 2.02 |
| 2.50 | All Liquid Discharge and Unplanned Release Sites in the 200 West Area Plus SALDS | L.52 |
| L.31 | SAC Results for Annual Vadose Zone Release of Uranium from All Liquid | 1.52 |
| L .51 | Discharge and Unplanned Release Sites in the 200 East Area | L.53 |
| L.32 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Uranium from | L.J. |
| L.32 | All Liquid Discharge and Unplanned Release Sites in the 200 East Area | L.53 |
| L.33 | SAC Results for Annual Vadose Zone Release of Uranium from All Liquid | L.J. |
| L.33 | Discharge and Unplanned Release Sites in the 200 West Area Plus SALDS | L.54 |
| L.34 | SAC Results for Cumulative Vadose Zone Release of Uranium from All Liquid | L.J- |
| L.34 | • | L.54 |
| I 25 | Discharge and Unplanned Release Sites in the 200 West Area Plus SALDS | L.32 |
| L.35 | SAC Results for Annual Vadose Zone Release of Iodine-129 from All Liquid Discharge | T 55 |
| 1 26 | and Unplanned Release Sites in the 200 East Area | L.55 |
| L.36 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Iodine-129 from | T ~ |
| 1 27 | All Liquid Discharge and Unplanned Release Sites in the 200 East Area | L.55 |
| L.37 | SAC Results for Annual Vadose Zone Release of Iodine-129 from All Liquid Discharge | T ~ |
| T 20 | and Unplanned Release Sites in the 200 West Area plus SALDS | L.56 |
| L.38 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Iodine-129 from | |
| | All Liquid Discharge and Unplanned Release Sites in the 200 West Area Plus SALDS | L.56 |
| L.39 | SAC Results for Annual Vadose Zone Release of Technetium-99 from All Other | . |
| * 40 | Sites in the 200 East Area | L.58 |
| L.40 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Technetium-99 from | |
| | All Other Sites in the 200 East Area | L.58 |
| L.41 | SAC Results for Annual Vadose Zone Release of Technetium-99 from All Other | |
| | Sites in the 200 West Area | L.59 |
| L.42 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Technetium-99 from | |
| | All Other Sites in the 200 West Area | L.59 |
| L.43 | SAC Results for Annual Vadose Zone Release of Technetium-99 from all Other | |
| | Sites Outside the 200 East and 200 West Areas | L.60 |
| L.44 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Technetium-99 from | |
| | All Other Sites Outside the 200 East and 200 West Areas | L.60 |
| L.45 | SAC Results for Annual Vadose Zone Release of Uranium from All Other Sites | |
| | in the 200 East Area | L.61 |
| L.46 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Uranium from All | |
| | Other Sites in the 200 East Area | L.61 |

| L.4/ | SAC Results for Annual Vadose Zone Release of Uranium from All Other Sites | |
|------|--|------|
| | in the 200 West Area | L.62 |
| L.48 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Uranium from All | |
| | Other Sites in the 200 West Area | L.62 |
| L.49 | SAC Results for Annual Vadose Zone Release of Uranium from All Other Sites | |
| | Outside the 200 East and 200 West Areas | L.63 |
| L.50 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Uranium from All | |
| | Other Sites Outside the 200 East and 200 West Areas | L.63 |
| L.51 | SAC Results for Annual Vadose Zone Release of Iodine-129 from All Other Sites in the 200 East Area | L.64 |
| L.52 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Iodine-129 from | |
| | All Other Sites in the 200 East Area | L.64 |
| L.53 | SAC Results for Annual Vadose Zone Release of Iodine-129 from All Other Sites in | |
| | the 200 West Area | L.65 |
| L.54 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Iodine-129 from | |
| | All Other Sites in the 200 West Area | L.65 |
| L.55 | SAC Results for Annual Vadose Zone Release of Iodine-129 from All Other Sites | |
| | Outside the 200 East and 200 West Areas. | L.66 |
| L.56 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Iodine-129 from | |
| | All Other Sites Outside the 200 East and 200 West Areas. | L.66 |
| L.57 | SAC Results for Annual Vadose Zone Release of Technetium-99 from ERDF | L.67 |
| L.58 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Technetium-99 | |
| | from ERDF | L.67 |
| L.59 | SAC Results for Annual Vadose Zone Release of Uranium from the ERDF | L.68 |
| L.60 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Uranium | I (0 |
| T (1 | from the ERDF | L.68 |
| L.61 | SAC Results for Annual Vadose Zone Release of Iodine-129 from ERDF | L.69 |
| L.62 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Iodine-129 | L.69 |
| 1 62 | from ERDF | L.05 |
| L.63 | | L.70 |
| I 61 | Commercial Low-Level Radioactive Waste Disposal Site | L./(|
| L.64 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Technetium-99 | 1 70 |
| I 65 | from the Commercial Low-Level Radioactive Waste Disposal Site | L.70 |
| L.65 | SAC Results for Annual Vadose Zone Release of Uranium from the Commercial | T 71 |
| I ((| Low-Level Radioactive Waste Disposal Site | L.71 |
| L.66 | SAC Results for Cumulative (undecayed) Vadose Zone Release of Uranium from the | T 71 |
| I 47 | Commercial Low-Level Radioactive Waste Disposal Site | L.71 |
| L.67 | | L.72 |
| L.68 | Low-Level Radioactive Waste Disposal Site | L./2 |
| L.00 | | L.72 |
| L.69 | from the Commercial Low-Level Radioactive Waste Disposal | L./2 |
| L.09 | | |
| | Except ILAW, Melters, and Naval Reactors in Groundwater 1 km Downgradient | L.76 |
| L.70 | Northeasterly of the 200 West Area | L./C |
| L./U | Except ILAW, Melters, and Naval Reactors in Groundwater 1 km Downgradient | |
| | Northeasterly of the 200 West Area | 1 77 |
| | INDITUTE ASTELLY OF THE ZOO WEST ATEA | ⊥.// |

| L./1 | Hypothetical Drinking Water Dose from Iodine-129 from All Hanford Sources | |
|------|--|------|
| | Except ILAW, Melters, and Naval Reactors in Groundwater 1 km Downgradient | |
| | Northeasterly of the 200 West Area | L.77 |
| L.72 | Hypothetical Drinking Water Dose from Technetium-99 from All Hanford Sources | |
| | Except ILAW, Melters, and Naval Reactors in Groundwater 1 km Downgradient | |
| | Southeasterly of the 200 East Area | L.78 |
| L.73 | Hypothetical Drinking Water Dose from Uranium from All Hanford Sources | |
| | Except ILAW, Melters, and Naval Reactors in Groundwater 1 km Downgradient | |
| | Southeasterly of the 200 East Area | L.78 |
| L.74 | Hypothetical Drinking Water Dose from Iodine-129 from All Hanford Sources | |
| | Except ILAW, Melters, and Naval Reactors in Groundwater 1 km Downgradient | |
| | Southeasterly of the 200 East Area | L.79 |
| L.75 | Hypothetical Drinking Water Dose from Technetium-99 from All Hanford Sources | |
| | Except ILAW, Melters, and Naval Reactors in Groundwater 1 km Downgradient | |
| | Northwesterly of the 200 East Area | L.80 |
| L.76 | Hypothetical Drinking Water Dose from Uranium from All Hanford Sources | |
| | Except ILAW, Melters, and Naval Reactors in Groundwater 1 km Downgradient | |
| | Northwesterly of the 200 East Area | L.80 |
| L.77 | Hypothetical Drinking Water Dose from Iodine-129 from All Hanford Sources | |
| | Except ILAW, Melters, and Naval Reactors in Groundwater 1 km Downgradient | |
| | Northwesterly of the 200 East Area | L.81 |
| L.78 | Drinking Water Dose at the City of Richland Pumping Station from Technetium-99 | |
| | Due to All Hanford Sources Except ILAW, Melters, and Naval Reactors | L.82 |
| L.79 | Drinking Water Dose at the City of Richland Pumping Station from Uranium Due to | |
| * 00 | All Hanford Sources Except ILAW, Melters, and Naval Reactors | L.83 |
| L.80 | Drinking Water Dose at the City of Richland Pumping Station from Iodine-129 Due | |
| T 04 | to All Hanford Sources Except ILAW, Melters, and Naval Reactors | L.83 |
| L.81 | Hypothetical Drinking Water Dose from Technetium-99 from Hanford Sources | |
| T 00 | Including ILAW in Groundwater 1 km Downgradient Southeasterly of the 200 East Area. | L.85 |
| L.82 | Hypothetical Drinking Water Dose from Uranium from Hanford Sources Including | T 05 |
| 1 02 | ILAW in Groundwater 1 km Downgradient Southeasterly of the 200 East Area | L.85 |
| L.83 | Hypothetical Drinking Water Dose from Iodine-129 from Hanford Sources Including | I 00 |
| T 04 | ILAW in Groundwater 1 km Downgradient Southeasterly of the 200 East Area | L.86 |
| L.84 | Concentrations of Technetium-99, Iodine-129, and Uranium from All Hanford Sources | T 06 |
| T 05 | in Groundwater 1 km Downgradient Southeasterly of 200 East Area. | L.86 |
| L.85 | Concentrations of Technetium-99, Iodine-129, and Uranium from All Hanford Sources | 1 07 |
| 1 06 | in Groundwater 1 km Downgradient Southeasterly of 200 East Area. | L.87 |
| L.86 | Hypothetical Total Drinking Water Dose from All Hanford Sources and from Solid | L.87 |
| L.87 | Waste Contributions in Groundwater 1 km Downgradient Southeasterly of 200 East Area. | L.0/ |
| L.0/ | Annual Drinking Water Dose from Technetium-99 in the Columbia River at the City | 1 00 |
| T 00 | of Richland Pumping Station from Hanford Sources Including ILAW | L.90 |
| L.88 | Annual Drinking Water Dose from Iodine-129 in the Columbia River at the City | 1 00 |
| L.89 | of Richland Pumping Station from Hanford Sources Including ILAW Annual Drinking Water Dose from Uranium in the Columbia River at the City | L.90 |
| 止.07 | · · · · · · · · · · · · · · · · · · · | L.91 |
| L.90 | of Richland Pumping Station from Hanford Sources Including ILAW Concentration of Technetium-99, Iodine-129, and Uranium in the Columbia River | L.71 |
| L.7U | at the City of Richland Pumping Station | L.91 |
| | at the vary of Newhall Editions Mailer | |

| L.91 | Drinking Water Dose from Technetium-99, Iodine-129, and Uranium in the Columbia | |
|------------|---|------------|
| 1 00 | River at the City of Richland Pumping Station. | L.92 |
| L.92 | Total Drinking Water Dose from All Hanford Sources and the Solid Waste Contribution in the Columbia River at the City of Richland Pumping Station | L.92 |
| M.1 | Hypothetical Annual Dose from Drinking Water Containing Maximum Combined Concentrations of Radionuclides in Groundwater at 1 km Downgradient from the | MO |
| M.2 | Hypothetical Annual Dose from Drinking Water Containing Maximum Combined Concentrations of Radionuclides in Groundwater 1 km Downgradient Northwest of the | M.2 M.2 |
| M.3 | Hypothetical Annual Dose from Drinking Water Containing Maximum Combined Concentrations of Radionuclides in Groundwater Near the Columbia River as a Function | M.3 |
| M.4 | Annual Dose to a Hypothetical Resident Gardener at Various Times over | M.4 |
| M.5 | Annual Dose to a Hypothetical Resident Gardener at Various Times over 10,000 Years Using Water from a Well 1 km Downgradient Northwest of the | M.4 |
| M.6 | Annual Dose to a Hypothetical Resident Gardener at Various Times over | M.5 |
| M.7 | Annual Dose to a Hypothetical Resident Gardener with a Sauna/Sweat Lodge Scenario at Various Times over 10,000 Years Using Water from a Well 1 km Downgradient from | |
| M.8 | the 200 West Area | M.5 |
| M.9 | Northwest of the 200 East Area | M.6 |
| | Columbia River | M.6 |
| N.1 | Hanford's Land-Use Plan | N.12 |
| O.1 | | |
| | Tables | |
| A.1 | Individuals, Organizations, and Agencies that Commented on the Scoping Phase of the ILAW SEIS | A.2 |
| A.2 A.3 | ILAW Disposal SEIS – Public Scoping Comments and Responses Individuals, Organizations, and Agencies Commenting on the Scoping Phase of the | A.3 |
| | HSW EIS | A.21 |
| B.1 | Assumptions for Management of Low-Level Waste | B.4 |
| B.2 | Assumptions for Management of Mixed Low-Level Waste | B.5 |

| B.3 | Assumptions for Management of Transuranic Waste | B.7 |
|-------|--|------|
| B.4 | Assumptions for Management of Waste Treatment Plant Wastes | B.9 |
| B.5 | Low-Level Waste Hanford Only Volumes | B.10 |
| B.6 | Low-Level Waste Lower Bound Volumes | B.10 |
| B.7 | Low-Level Waste Upper Bound Volumes | B.10 |
| B.8 | Mixed Low-Level Waste Hanford Only Volumes | B.11 |
| B.9 | Mixed Low-Level Waste Lower Bound Volumes | B.11 |
| B.10 | Mixed Low-Level Waste Upper Bound Volumes | B.12 |
| B.11 | Transuranic Waste Hanford Only Volumes | B.12 |
| B.12 | Transuranic Waste Lower Bound Volumes | B.13 |
| B.13 | Transuranic Waste Upper Bound Volumes | B.13 |
| B.14 | Waste Treatment Plant Waste Volumes | B.14 |
| B.15 | Inventory of Long-Lived Mobile Radionuclides in HSW for the Various | |
| | Alternative Groups, Ci | B.16 |
| B.16 | Inventory of Long-Lived Mobile Radionuclides in HSW for Alternative Group A, Ci | B.17 |
| B.17 | Inventory of Long-Lived Mobile Radionuclides in HSW for Alternative Group B, Ci | B.21 |
| B.18 | Inventory of Long-Lived Mobile Radionuclides in HSW for Alternative Group C, Ci | B.25 |
| B.19 | Inventory of Long-Lived Mobile Radionuclides in HSW for Alternative Groups D ₁ | |
| | and D ₂ , Ci | B.29 |
| B.20 | Inventory of Long-Lived Mobile Radionuclides in HSW for Alternative Group D ₃ , Ci | B.33 |
| B.21 | Inventory of Long-Lived Mobile Radionuclides in HSW for Alternative Groups E ₁ | |
| | and E ₂ , Ci | B.37 |
| B.22 | Inventory of Long-Lived Mobile Radionuclides in HSW for Alternative Group E ₃ , Ci | B.41 |
| B.23 | Inventory of Long-Lived Mobile Radionuclides in HSW for the No Action | |
| | Alternative, Ci | B.45 |
| B.24a | Inventory of MLLW as Soil and Grouted-Equivalent Fractions for Alternative Groups A, C, D, and E, Ci | B.46 |
| B.24b | Inventory of MLLW as Soil and Grouted-Equivalent Fractions for Alternative | |
| | Group B, Ci. | B.47 |
| B.24c | Inventory of MLLW as Soil and Grouted-Equivalent Fractions for the No Action | |
| | Alternative, Ci | B.48 |
| B.25 | Identification of Flowsheets | B.49 |
| B.26 | Matrix of Disposal Options for Alternative Groups D and E | B.81 |
| | • | |
| C.1. | Comparison of Waste Management Programmatic Environmental Impact Statement, | |
| | Accelerated Cleanup: Paths to Closure, and HSW EIS Waste Volumes | C.3 |
| C.2 | Hanford Only Volume for Low-Level Waste | C.5 |
| C.3 | Lower Bound Volume for Low-Level Waste | C.6 |
| C.4 | Upper Bound Volume for Low-Level Waste | C.7 |
| C.5 | Hanford Only Volume for Mixed Low-Level Waste | C.7 |
| C.6 | Lower Bound Volume for Mixed Low-Level Waste | C.8 |
| C.7 | Upper Bound Volume for Mixed Low-Level Waste | C.9 |
| C.8 | Hanford Only Waste Volumes for Transuranic Waste | C.10 |
| C.9 | Lower Bound Waste Volumes for Transuranic Waste | C.10 |
| C.10 | Upper Bound Waste Volumes for Transuranic Waste | C.11 |
| C.11 | Estimated Volumes of WTP Waste Streams through 2046 | C.11 |

| E.1 | 200 East and 200 West Area Emissions: Dispersion Factors Used to Determine | |
|------|--|------|
| | Maximum Air Quality Impacts to the Public | E.4 |
| E.2 | Area C Borrow Pit Emissions: Location and Dispersion Factors Used to Determine | |
| | Maximum Air Quality Impacts | E.5 |
| E.3 | Timeline of Alternative Group A Activities Resulting in Criteria Pollutant Emissions | E.7 |
| E.4 | Timeline of Alternative Group B Activities Resulting in Criteria Pollutant Emissions | E.8 |
| E.5 | Timeline of Alternative Group C Activities Resulting in Criteria Pollutant Emissions | E.9 |
| E.6 | Timeline of Alternative Group D Activities Resulting in Criteria Pollutant Emissions | E.10 |
| E.7 | Timeline of Alternative Group E Activities Resulting in Criteria Pollutant Emissions | E.11 |
| E.8 | Timeline of the No Action Alternative Resulting in Criteria Pollutant Emissions | E.12 |
| E.9 | Emission Factors for Criteria Pollutants | E.13 |
| E.10 | Average Diesel Fuel Consumption Rates | E.15 |
| E.11 | Average Propane Fuel Consumption Rates | E.17 |
| E.12 | Size of Disturbed Areas and Associated Durations for Various Activities/ | |
| | Alternatives | E.20 |
| E.13 | Maximum Air Quality Impacts to the Public from Major Activities with a Source | |
| | Location in the 200 West or 200 East Areas | E.22 |
| E.14 | Maximum Impacts from Any Single Activity Conducted in the 200 Areas | E.25 |
| E.15 | Maximum Air Quality Impacts to the Public from Activities with an Area C | |
| | Source Location | E.25 |
| E.16 | Clean Air Act Conformity Review for the Alternatives | E.26 |
| | | |
| F.1 | Summary of Waste Stream Concentration Tables | F.3 |
| F.2 | Stream 1: Low-Level Waste Category 1 | F.4 |
| F.3 | Stream 2: Low-Level Waste Category 3 | F.4 |
| F.4 | Streams 1 and 2: Low-Level Waste from Offsite Sources | F.5 |
| F.5 | Stream 2C2: Low-Level Waste Category 3 for T Plant Processing from Offsite Sources | F.6 |
| F.6 | Stream 4: TRU-RH Waste from Trenches | F.7 |
| F.7 | Stream 4: TRU-CH Waste from Trenches | F.7 |
| F.8 | Stream 5: TRU-CH Waste from Caissons | F.7 |
| F.9 | Stream 8: TRU – Commingled Polychlorinated Biphenyl Waste | F.8 |
| F.10 | Streams 9 and 10A: TRU – Newly Generated and Existing CH Standard and | |
| | Non-Standard Containers | F.8 |
| F.11 | Stream 10B: TRU – Newly Generated and Existing RH Waste | F.9 |
| F.12 | Stream 11: MLLW Treated and Ready for Disposal | F.9 |
| F.13 | Stream 12: RH and Non-Standard Packages | F.11 |
| F.14 | Stream 13: CH Organic and Inorganic Solids and Debris | F.11 |
| F.15 | Stream 14: Elemental Lead | F.13 |
| F.16 | Stream 15: Elemental Mercury | F.14 |
| F.17 | Stream 17: K Basin Sludge | F.14 |
| F.18 | Stream 18: MLLW Trench Leachate | F.15 |
| F.19 | Release Fraction Values for the WRAP | F.16 |
| F.20 | Release Fractions for Volatile Organic Compounds from the WRAP | F.17 |
| F.21 | Airborne Radionuclide Releases from the WRAP | F.18 |
| F.22 | Total Chemical Atmospheric Releases from the WRAP | F.19 |
| F.23 | Release Fraction Values for the 2706-T Facility in the T Plant Complex | F.19 |
| F.24 | Total Radionuclide Atmospheric Release from the T Plant Complex | F.20 |
| F.25 | Total Chemical Atmospheric Releases from the T Plant Complex | F.21 |

| F.26 | Release Fraction Values for the New Waste Processing Facility and the | |
|------|--|------|
| | Modified T Plant Complex | F.22 |
| F.27 | Total Radionuclide Atmospheric Release from the Modified T Plant Complex for | |
| | Alternative Group A | F.22 |
| F.28 | Total Chemical Atmospheric Releases from the Modified T Plant Complex for | |
| | Alternative Group A | F.23 |
| F.29 | Atmospheric Radionuclide Releases from the New Waste Processing Facility for | |
| | Alternative Group B | F.23 |
| F.30 | Total Chemical Atmospheric Releases from the New Waste Processing Facility for | |
| | Alternative Group B | F.24 |
| F.31 | Atmospheric Radionuclide Release from Pulse Drier Leachate Treatment— | |
| | Alternative Groups A and B | F.25 |
| F.32 | Atmospheric Radionuclide Release from Pulse Drier Leachate Treatment— | |
| | Alternative Groups C and D, and the No Action Alternative | F.26 |
| F.33 | Release Point Characteristics | F.27 |
| F.34 | Joint Frequency Distributions for the 200 Areas at 9.1-m Towers, 1983-1996 | |
| | Historical Data | F.28 |
| F.35 | Joint Frequency Distributions for the 200 Areas at 60-m Aboveground Level, | |
| | 1983-1996 Historical Data | F.29 |
| F.36 | Population Within 80 km of the 200 Areas | F.32 |
| F.37 | Industrial Scenario Exposure Pathways | F.34 |
| F.38 | Industrial Scenario Parameter Values | F.34 |
| F.39 | Resident Gardener Scenario Exposure Pathways | F.35 |
| F.40 | Resident Gardener Scenario Parameter Values | F.36 |
| F.41 | Volatilization Half-Times for Soil | F.40 |
| F.42 | Soil Classification Composition | F.41 |
| F.43 | Soil-Related Properties of Hazardous Chemicals | F.42 |
| F.44 | Distribution Coefficients of Radionuclides and Inorganic Chemicals | F.43 |
| F.45 | Summary of Basis for Health Consequences from Radiological Exposures from | |
| | Federal Guidance Report 13 | F.44 |
| F.46 | Basis for Health Consequences from Radiological Exposures | F.45 |
| F.47 | Ratios of Dose and Risk to Children over Dose and Risk to Adults from 1-Year | |
| | Ingestion of Contaminated Drinking Water | F.46 |
| F.48 | Peak Radionuclide Concentrations in Disposal Facilities | F.52 |
| F.49 | Dose-Rate-per-Unit Waste Concentration Factors for the Post-Well Drilling | |
| | Scenario, Time Since Year 2046 | F.54 |
| F.50 | Dose-Rate-per-Unit Waste Concentration Factors for the Excavation Scenario, | |
| | Time Since Year 2046 | F.55 |
| F.51 | Content of Figures for Groundwater Analysis Results | F.56 |
| F.52 | Exposure Scenario Dose Factors for Use of Groundwater | F.94 |
| F.53 | Content of Tables for Groundwater Analysis Results | F.95 |
| F.54 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener | |
| | at the 200 East Area 1-km Well from Radionuclides in the Groundwater over | |
| | 10,000 Years for Alternative Group A, Hanford Only Waste Volume | F.96 |
| F.55 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener | |
| | at the 200 East Area 1-km Well from Radionuclides in the Groundwater over | |
| | 10.000 Years for Alternative Group A. Lower Bound Waste Volume | F.97 |

| F.50 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener | |
|------|--|-------|
| | at the 200 East Area 1-km Well from Radionuclides in the Groundwater over | |
| | 10,000 Years for Alternative Group A, Upper Bound Waste Volume | F.98 |
| F.57 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener | |
| | at the 200 West Area 1-km Well from Radionuclides in the Groundwater over | |
| | 10,000 Years for Alternative Group A, Hanford Only Waste Volume | F.99 |
| F.58 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener | |
| | at the 200 West Area 1-km Well from Radionuclides in the Groundwater over | |
| | 10,000 Years for Alternative Group A, Lower Bound Waste Volume | F 100 |
| F.59 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener | 1.100 |
| 1.57 | at the 200 West Area 1-km Well from Radionuclides in the Groundwater over | |
| | 10,000 Years for Alternative Group A, Upper Bound Waste Volume | F 101 |
| F.60 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener | 1.101 |
| 1.00 | at the Columbia River Well from Radionuclides in the Groundwater over | |
| | | E 100 |
| Г (1 | 10,000 Years for Alternative Group A, Hanford Only Waste Volume | F.102 |
| F.61 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener | |
| | at the Columbia River Well from Radionuclides in the Groundwater over | E 100 |
| | 10,000 Years for Alternative Group A, Lower Bound Waste Volume | F.103 |
| F.62 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener | |
| | at the Columbia River Well from Radionuclides in the Groundwater over | |
| | 10,000 Years for Alternative Group A, Upper Bound Waste Volume | F.104 |
| F.63 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener | |
| | at the 200 East Area 1-km Well from Radionuclides in the Groundwater over | |
| | 10,000 Years for Alternative Group B, Hanford Only Waste Volume | F.105 |
| F.64 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener | |
| | at the 200 East Area 1-km Well from Radionuclides in the Groundwater over | |
| | 10,000 Years for Alternative Group B, Lower Bound Waste Volume | F.106 |
| F.65 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener | |
| | at the 200 East Area 1-km Well from Radionuclides in the Groundwater over | |
| | 10,000 Years for Alternative Group B, Upper Bound Waste Volume | F.107 |
| F.66 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener | |
| | at the 200 West Area 1-km Well from Radionuclides in the Groundwater over | |
| | 10,000 Years for Alternative Group B, Hanford Only Waste Volume | F.108 |
| F.67 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener | |
| | at the 200 West Area 1-km Well from Radionuclides in the Groundwater over | |
| | 10,000 Years for Alternative Group B, Lower Bound Waste Volume | F 109 |
| F.68 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener | 1.103 |
| 1.00 | at the 200 West Area 1-km Well from Radionuclides in the Groundwater over | |
| | 10,000 Years for Alternative Group B, Upper Bound Waste Volume | F 110 |
| F.69 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener | 1.11(|
| 1.07 | at the Columbia River Well from Radionuclides in the Groundwater over | |
| | 10,000 Years for Alternative Group B, Hanford Only Waste Volume | E 111 |
| E 70 | | Г.111 |
| F.70 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener | |
| | at the Columbia River Well from Radionuclides in the Groundwater over | D 110 |
| F 71 | 10,000 Years for Alternative Group B, Lower Bound Waste Volume | F.112 |
| F.71 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | Columbia River Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group B, Upper Bound Waste Volume | F.113 |

| F.72 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
|------|--|-------|
| | Alternative Group C, Hanford Only Waste Volume | F.114 |
| F.73 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group C, Lower Bound Waste Volume | F.115 |
| F.74 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group C, Upper Bound Waste Volume | F.116 |
| F.75 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | 200 West Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group C, Hanford Only Waste Volume | F.117 |
| F.76 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | 200 West Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group C, Lower Bound Waste Volume | F.118 |
| F.77 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | 200 West Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group C, Upper Bound Waste Volume | F.119 |
| F.78 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | Columbia River Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group C, Hanford Only Waste Volume | F.120 |
| F.79 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | Columbia River Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group C, Lower Bound Waste Volume | F.121 |
| F.80 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | Columbia River Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group C, Upper Bound Waste Volumes | F.122 |
| F.81 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | E 100 |
| F 02 | Alternative Group D ₁ , Hanford Only Waste Volume | F.123 |
| F.82 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | E 104 |
| E 02 | Alternative Group D ₁ , Lower Bound Waste Volume | F.124 |
| F.83 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | E 105 |
| F.84 | Alternative Group D ₁ , Upper Bound Waste Volume | F.125 |
| Г.04 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | 200 West Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | E 126 |
| F.85 | Alternative Group D ₁ , Hanford Only Waste Volume | F.126 |
| г.оэ | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the 200 West Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group D ₁ , Lower Bound Waste Volume | F.127 |
| F.86 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | 1.12/ |
| 1.00 | 200 West Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group D ₁ , Upper Bound Waste Volume | F.128 |
| F.87 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | 1.120 |
| 1.07 | Columbia River Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group D ₁ . Hanford Only Waste Volume | F.129 |

| F.88 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
|-------|--|-------|
| | Columbia River Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group D ₁ , Lower Bound Waste Volume | F.130 |
| F.89 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | Columbia River Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group D ₁ , Upper Bound Waste Volume | F.131 |
| F.90 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | 1.13 |
| 1.70 | - · · · · · · · · · · · · · · · · · · · | |
| | 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | E 120 |
| T 0.1 | Alternative Group D ₂ , Hanford Only Waste Volume | F.132 |
| F.91 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group D ₂ , Lower Bound Waste Volume | F.133 |
| F.92 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group D ₂ , Upper Bound Waste Volume | F.134 |
| F.93 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | 200 West Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group D ₂ , Hanford Only Waste Volume | F.135 |
| F.94 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | 200 West Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group D ₂ , Lower Bound Waste Volume | F.136 |
| F.95 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | 1.150 |
| 1.75 | 200 West Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group D ₂ , Upper Bound Waste Volume | F.137 |
| E 06 | | 1.13 |
| F.96 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | Columbia River Well from Radionuclides in the Groundwater over 10,000 Years for | E 100 |
| F.05 | Alternative Group D ₂ , Hanford Only Waste Volume | F.138 |
| F.97 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | Columbia River Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group D ₂ , Lower Bound Waste Volume | F.139 |
| F.98 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | Columbia River Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group D ₂ , Upper Bound Waste Volume | F.140 |
| F.99 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group D ₃ , Hanford Only Waste Volume | F.141 |
| F.100 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group D ₃ , Lower Bound Waste Volume | F.142 |
| F.101 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| 1.101 | 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group D ₃ , Upper Bound Waste Volume | F.143 |
| F.102 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | 1.17. |
| 1.102 | 200 West Area and 200 ERDF Site 1-km Wells from Radionuclides in the Groundwater | |
| | | E 144 |
| E 102 | over 10,000 Years for Alternative Group D ₃ , Hanford Only Waste Volume | F.144 |
| F.103 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | 200 West Area and 200 ERDF Site 1-km Wells from Radionuclides in the Groundwater | T 14 |
| | over 10,000 Years for Alternative Group D ₃ , Lower Bound Waste Volume | F.145 |

| F.104 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
|-------|--|-------|
| | 200 West Area and 200 ERDF Site 1-km Wells from Radionuclides in the Groundwater | |
| | over 10,000 Years for Alternative Group D ₃ , Upper Bound Waste Volume | F.146 |
| F.105 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | Columbia River Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group D ₃ , Hanford Only Waste Volume | F.147 |
| F.106 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | 1.1.7 |
| 1.100 | Columbia River Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | | F.148 |
| F 107 | Alternative Group D ₃ , Lower Bound Waste Volume | F.148 |
| F.107 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | Columbia River Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group D ₃ , Upper Bound Waste Volume | F.149 |
| F.108 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group E ₁ , Hanford Only Waste Volume | F.150 |
| F.109 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group E ₁ , Lower Bound Waste Volume | F.151 |
| F.110 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group E ₁ , Upper Bound Waste Volume | F.152 |
| F.111 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | 1110- |
| | 200 West Area and 200 ERDF Site 1-km Wells from Radionuclides in the Groundwater | |
| | over 10,000 Years for Alternative Group E ₁ , Hanford Only Waste Volume | F.153 |
| F.112 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | 1.133 |
| 1.112 | 200 West Area and 200 ERDF Site 1-km Wells from Radionuclides in the Groundwater | |
| | | F.154 |
| E 112 | over 10,000 Years for Alternative Group E ₁ , Lower Bound Waste Volume | Г.134 |
| F.113 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | 200 West Area and 200 ERDF Site 1-km Wells from Radionuclides in the Groundwater | D 155 |
| | over 10,000 Years for Alternative Group E ₁ , Upper Bound Waste Volume | F.155 |
| F.114 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | Columbia River Well from Radionuclides in the Groundwater over 10,000 Years for- | |
| | Alternative Group E ₁ , Hanford Only Waste Volume | F.156 |
| F.115 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | Columbia River Well from Radionuclides in the Groundwater Over 10,000 Years for | |
| | Alternative Group E ₁ , Lower Bound Waste Volume | F.157 |
| F.116 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | Columbia River Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group E ₁ , Upper Bound Waste Volume | F.158 |
| F.117 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group E ₂ , Hanford Only Waste Volume | F.159 |
| F.118 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | 1.10) |
| 1.110 | 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group E ₂ , Lower Bound Waste Volume | F.160 |
| F.119 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | 1.100 |
| 1,117 | 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group F. Upper Round Waste Volume | F 161 |
| | | |

| F.120 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
|-------|--|-------|
| | 200 West Area and 200 ERDF Site 1-km Wells from Radionuclides in the Groundwater | E 160 |
| F.121 | over 10,000 Years for Alternative Group E ₂ , Hanford Only Waste Volume | F.162 |
| Γ.121 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the 200 West Area and 200 ERDF Site 1-km Well from Radionuclides in the Groundwater | |
| | over 10,000 Years for Alternative Group E ₂ , Lower Bound Waste Volume | F.163 |
| F.122 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | Г.103 |
| 1.122 | 200 West Area and 200 ERDF Site 1-km Wells from Radionuclides in the Groundwater | |
| | | F.164 |
| E 122 | over 10,000 Years for Alternative Group E ₂ , Upper Bound Waste Volume | Г.102 |
| F.123 | Columbia River Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group E ₂ , Hanford Only Waste Volume | F.165 |
| E 124 | • | F.10. |
| F.124 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | |
| | Columbia River Well from Radionuclides in the Groundwater over 10,000 Years for | F.166 |
| E 125 | Alternative Group E ₂ , Lower Bound Waste Volume | F.100 |
| F.125 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the Columbia River Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group E ₂ , Upper Bound Waste Volume | F.167 |
| F.126 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | 1.10 |
| 1.120 | 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group E ₃ , Hanford Only Waste Volume | F.168 |
| F.127 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | Г.100 |
| 1.12/ | 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group E ₃ , Lower Bound Waste Volume | F.169 |
| F.128 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | 1.105 |
| Г.120 | 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group E ₃ , Upper Bound Waste Volume | F.170 |
| F.129 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | 1.170 |
| 1.129 | 200 West Area and 200 ERDF Site 1-km Wells from Radionuclides in the Groundwater | |
| | over 10,000 Years for Alternative Group E ₃ , Hanford Only Waste Volume | F.171 |
| F.130 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | 1.1/1 |
| 1.130 | 200 West Area and 200 ERDF Site 1-km Wells from Radionuclides in the Groundwater | |
| | over 10,000 Years for Alternative Group E ₃ , Lower Bound Waste Volume | F.172 |
| F.131 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | 1.1/2 |
| 1.131 | 200 West Area and 200 ERDF Site 1-km Wells from Radionuclides in the Groundwater | |
| | over 10,000 Years for Alternative Group E ₃ , Upper Bound Waste Volume | F 173 |
| F.132 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | 1.1/5 |
| 1.132 | Columbia River Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group E ₃ , Hanford Only Waste Volume | F.174 |
| F.133 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | 1.17- |
| 1.133 | Columbia River Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group E ₃ , Lower Bound Waste Volume | F.175 |
| F.134 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | 1.1/5 |
| 1.154 | Columbia River Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | Alternative Group E ₃ , Upper Bound Waste Volume | F.176 |
| F.135 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the | 1.1/(|
| 1.133 | 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| | the No Action Alternative, Hanford Only Waste Volume | F.177 |
| | | / / |

| F.136 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the 200 East Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | 7.450 |
|----------------|--|---------|
| F.137 | the No Action Alternative, Lower Bound Waste Volume | F.178 |
| | the No Action Alternative, Hanford Only Waste Volume | F.179 |
| F.138 | Potential Individual Human Health Impacts to a Hypothetical Resident Gardener at the 200 West Area 1-km Well from Radionuclides in the Groundwater over 10,000 Years for | |
| F.139 | the No Action Alternative, Lower Bound Waste Volume | F.180 |
| F.140 | the No Action Alternative, Hanford Only Waste Volume | F.181 |
| | the No Action Alternative, Lower Bound Waste Volume | F.182 |
| F.141 | West Valley TRU Radionuclide Concentrations | |
| F.142 F.143 | Occupational Exposure from Processing West Valley TRU Wastes at Hanford | F.184 |
| | Releases of Radionuclides – West Valley TRU Waste Only | F.185 |
| G.1 | Constituents Categorized by Mobility (K _d) Classes | G.10 |
| G.2 | Assumed Distribution of LLBG Areas of Previously Disposed of LLW, | 0.10 |
| 0.2 | Cat 1 LLW, Cat 3 LLW, MLLW, and Melters in the 200 East and 200 West Areas | |
| | by Alternative Group | G.13 |
| G.3 | Summary of Waste Depth and Infiltration Rates Used in the Soil-Debris Release Model | G.15 |
| G.4 | Geologic Well Logs for the Vadose Zone Model | G.22 |
| G.5 | Sediment Types and Unsaturated Flow Model Parameters Used | |
| | in the Composite Analysis | G.23 |
| G.6 | Major Hydrogeologic Units Used in the Sitewide Three-Dimensional Model | G.27 |
| G.7 | Predicted Peak Concentrations of Key Constituents from Wastes Disposed of | |
| | Before 2008 at a 1-km Line of Analysis, All Action Alternatives | G.78 |
| G.8 | Predicted Peak Concentrations of Key Constituents from Wastes Disposed of | |
| | Before 2008 at a Line of Analysis Near the Columbia River, All Action Alternatives | G.84 |
| G.9 | Predicted Peak River Flux of Key Constituents from Wastes Disposed of Before | |
| | 2008 at a Line of Analysis Near the Columbia River, All Action Alternatives | G.90 |
| G.10 | Predicted Peak Concentrations of Key Constituents Disposed of After 2007 at a | |
| | 1-km Line of Analysis, Alternative Group A | G.96 |
| G.11 | Predicted Peak Concentrations of Key Constituents Disposed of After 2007 at a | |
| | Line of Analysis Near the Columbia River, Alternative Group A | G.99 |
| G.12 | Predicted Peak River Flux of Key Constituents Disposed of After 2007 at a | |
| | Line of Analysis to the Columbia River, Alternative Group A | G.102 |
| G.13 | Predicted peak Concentrations of Key Constituents Disposed of After 2007 at a | |
| | 1-km Line of Analysis, Alternative Group B | G.105 |
| G.14 | Predicted Peak Concentrations of Key Constituents Disposed of After 2007 at a | |
| ~ | Line of Analysis Near the Columbia River, Alternative Group B | G.108 |
| G.15 | Predicted Peak River Flux of Key Constituents Disposed of After 2007 at a | |
| | Line of Analysis to the Columbia River. Alternative Group B | G.111 |

| G.10 | Predicted Peak Concentrations of Key Constituents Disposed of After 2007 at a | |
|-------|---|--------------|
| | 1-km Line of Analysis, Alternative Group C | G.114 |
| G.17 | Predicted Peak Concentrations of Key Constituents Disposed of After 2007 at a | |
| | Line of Analysis Near the Columbia River, Alternative Group C | G.117 |
| G.18 | Predicted Peak River Flux of Key Constituents Disposed of After 2007 at a | |
| | Line of Analysis to the Columbia River, Alternative Group C | G.120 |
| G.19 | Predicted Peak Concentrations of Key Constituents Disposed of After 2007 at a | |
| | 1-km Line of Analysis, Alternative Group D ₁ | G.123 |
| G.20 | Predicted Peak Concentrations of Key Constituents Disposed of After 2007 at a | |
| | Line of Analysis Near the Columbia River, Alternative Group D ₁ | G.126 |
| G.21 | Predicted Peak River Flux of Key Constituents Disposed of After 2007 at a | |
| | Line of Analysis to the Columbia River, Alternative Group D ₁ | G.129 |
| G.22 | Predicted Peak Concentrations of Key Constituents Disposed of After 2007 at a | |
| | 1-km Line of Analysis, Alternative Group D ₂ | G.132 |
| G.23 | Predicted Peak Concentrations of Key Constituents Disposed of After 2007 at a | 0.102 |
| 0.20 | Line of Analysis Near the Columbia River, Alternative Group D ₂ | G 135 |
| G.24 | Predicted Peak River Flux of Key Constituents Disposed of After 2007 at a | 0.100 |
| 0.2 . | Line of Analysis to the Columbia River, Alternative Group D ₂ | G 138 |
| G.25 | Predicted Peak Concentrations of Key Constituents Disposed of After 2007 at a | 0.150 |
| 0.20 | 1-km Line of Analysis, Alternative Group D ₃ | G 141 |
| G.26 | Predicted Peak Concentrations of Key Constituents Disposed of After 2007 at a | 0.111 |
| 0.20 | Line of Analysis Near the Columbia River, Alternative Group D ₃ | G 144 |
| G.27 | Predicted Peak River Flux of Key Constituents Disposed of After 2007 at a | 0.144 |
| 0.27 | Line of Analysis to the Columbia River, Alternative Group D ₃ | G 147 |
| G.28 | Predicted Peak Concentrations of Key Constituents Disposed of After 2007 at a | 0.147 |
| 0.20 | 1-km Line of Analysis, Alternative Group E ₁ | G 150 |
| G.29 | Predicted Peak Concentrations of Key Constituents Disposed of After 2007 at a | 0.150 |
| 0.27 | Line of Analysis Near the Columbia River, Alternative Group E ₁ | G 153 |
| G.30 | Predicted Peak River Flux of Key Constituents Disposed of After 2007 at a | 0.133 |
| 0.50 | Line of Analysis to the Columbia River, Alternative Group E ₁ | G 156 |
| G.31 | Predicted Peak Concentrations of Key Constituents Disposed of After 2007 at a | 0.150 |
| 0.51 | 1-km Line of Analysis, Alternative Group E ₂ | G 150 |
| G.32 | Predicted Peak Concentrations of Key Constituents Disposed of After 2007 at a | U.139 |
| 0.32 | Line of Analysis Near the Columbia River, Alternative Group E ₂ | C 162 |
| G.33 | | G.102 |
| G.33 | Predicted Peak River Flux of Key Constituents Disposed of After 2007 at a | C 165 |
| C 24 | Line of Analysis to the Columbia River, Alternative Group E ₂ | G.103 |
| G.34 | Predicted Peak Concentrations of Key Constituents Disposed of After 2007 at a | C 160 |
| C 25 | 1-km Line of Analysis, Alternative Group E ₃ Predicted Peak Concentrations of Key Constituents Disposed of After 2007 at a | G.108 |
| G.35 | · · · · · · · · · · · · · · · · · · · | C 171 |
| 0.26 | Line of Analysis Near the Columbia River, Alternative Group E ₃ | G.1/1 |
| G.36 | Predicted Peak River Flux of Key Constituents Disposed of After 2007 at a | C 174 |
| C 27 | Line of Analysis to the Columbia River, Alternative E ₃ | G.1/4 |
| G.37 | Predicted Peak Concentrations of Key Constituents at a 1-km Line of Analysis, | G 155 |
| G 20 | No Action Alternative. | G.17/ |
| G.38 | Predicted Peak Concentrations of Key Constituents at a Line of Analysis | a 10: |
| G 60 | Near the Columbia River, No Action Alternative | G.181 |
| G.39 | Predicted Peak River Flux of Key Constituents at a Line of Analysis Near the | a 10= |
| | Columbia River, No Action Alternative | G.185 |

| G.40 | Drinking Water Doses (mrem/yr) | G.267 |
|-------|--|-------|
| G.41 | Well Intercept Factors at LOAs Downgradient from the ILAW Disposal Facility Sited | |
| | Near the PUREX Plant and Alternative Locations | G.271 |
| G.42 | Predicted Peak Concentrations of Key Constituents from Waste Disposed of | |
| | Before 2008 at Aggregate LLW Management Area Boundaries, Alternative | |
| | Groups D_1 , D_2 , and D_3 | G.284 |
| G.43 | Predicted Peak Concentrations of Key Constituents from Wastes Disposed of | 0.20 |
| 0.15 | After 2007 at Aggregate LLW Management Area Boundaries, Alternative Group D ₁ | G.288 |
| G.44 | Predicted Peak Concentrations of Key Constituents by from Wastes Disposed of | 0.200 |
| 0.44 | After 2007 at Aggregate LLW Management Area Boundaries, Alternative Group D ₂ | G.290 |
| C 15 | | G.290 |
| G.45 | Predicted Peak Concentrations of Key Constituents for Wastes Disposed of After 2007 | C 202 |
| G 46 | at Aggregate LLW Management Area Boundaries, Alternative Group D ₃ | G.292 |
| G.46 | Sum of MCL Fractions and Drinking Water Dose from Maximum Potential | |
| | Concentrations for Technetium-99 and Iodine-129 for Waste Buried Before 2008 | |
| | at Facility Boundaries | G.298 |
| G.47 | Sums of MCL Fractions and Drinking Water Doses from Maximum Potential | |
| | Concentrations for Technetium-99 and Iodine-129 for Waste Buried After 2007 | |
| | at Facility Boundaries | G.300 |
| G.48 | Estimated Inventories of Selected Hazardous Chemicals Potentially Disposed of in | |
| | HSW LLBGs Between 1962 and 1987 | G.301 |
| G.49 | Constituents Categorized by Mobility Classes | G.304 |
| G.50 | Degradation Rates of Selected Organic Chemicals Hypothetically | |
| | Associated with Waste Disposed of Before 1988 | G 305 |
| G.51 | Degradation Rates Due to Volatilization of Selected Organic Chemicals | 0.505 |
| 0.51 | Hypothetically Associated with Waste Disposed of Before 1988 Using | |
| | Methods by Strenge and Peterson | G.305 |
| C 52 | Predicted Peak Concentrations of Selected Hazardous Chemical | 0.505 |
| G.52 | | C 200 |
| | Within Waste Disposed of Before 1988 | G.300 |
| H.1 | Conoral Chinning Decemptors for HCW FIC Colid Wests Chinments | нο |
| | General Shipping Parameters for HSW EIS Solid Waste Shipments | |
| H.2 | Shipping Data for Alternative Group A, Hanford Only Waste Volume | |
| H.3 | Shipping Data for Alternative Group B, Hanford Only Waste Volume | |
| H.4 | Onsite Shipping Data for the No Action Alternative | |
| H.5 | Offsite Shipment Volumes and Shipment Projections | |
| H.6 | Shipping Capacities Used to Estimate Offsite Shipments | |
| H.7 | Route Characteristics Data for Offsite Shipments | H.21 |
| H.8 | RADTRAN 5 Accident Parameters for Onsite Truck Shipments | H.24 |
| H.9 | Radionuclide Concentrations Used to Calculate Per-Shipment Inventories | |
| | for Onsite Shipments | H.26 |
| H.10 | Radionuclide Inventories for Offsite Shipments | H.27 |
| H.11 | Hazardous Chemical Inventories in Various Waste Types | H.29 |
| H.12 | Transportation Impacts of Alternative Group A – Hanford Only Waste Volume, | |
| | Number of Fatalities | H.32 |
| H.13 | Transportation Impacts of Alternative Group B – Hanford Only Waste Volume, | |
| 11.13 | Number of Fatalities | H.35 |
| H.14 | Transportation Impacts for the No Action Alternative, Hanford Only Waste Volume, | 11.55 |
| 11.14 | Number of Fatalities | H.37 |
| H.15 | Summary of Impacts of Shipping Hanford Only Wastes for Each Alternative Group | H.40 |
|) | SUBBLIALY OF HUDAGIS OF SUBDIUS FLABBORD VIIIV WASIES TO FACE AUCHAUVE CHOID | 1141 |

| H.16 | Radiological Transportation Impacts for Offsite Shipments | H.41 |
|------------|--|------|
| H.17 | Summary of Radiological Transportation Impacts for Offsite Shipments by Waste Type | H.44 |
| H.18 | Non-Radiological Transportation Impacts for Offsite Shipments | H.45 |
| H.19 | Summary of Non-Radiological Impacts for Offsite Shipments by Waste Type (Fatalities) | H.47 |
| H.20 | Estimated Doses and Impacts to MEIs | H.48 |
| H.21 | Summary of Impacts of Maximum Credible Accidents from DOE (1997b) | H.49 |
| H.22 | Total Shipment Mileages by Shipment Origin | H.51 |
| H.23 | Latent Cancer Fatality Impacts by Shipment Origin | H.52 |
| H.24 | Non-Radiological Accident Impacts by Shipment Origin | H.53 |
| H.25 | Results of Sensitivity Study for Shifting TRU Waste Shipments from | |
| | California to New York | H.57 |
| H.26 | Potential Incremental Transportation Impacts if West Valley TRU Waste | |
| | were to be Shipped to Hanford | H.60 |
| H.27 | Total Potential HSW Transportation Impacts With and Without | |
| | West Valley TRU Waste Shipments | H.62 |
| H.28 | Impacts of Transporting Construction and Backfill Materials | H.64 |
| H.29 | Route Characteristics for Transport Within Washington and Oregon | H.68 |
| H.30 | Offsite Shipping Volumes Used for Oregon and Washington Impacts Calculations | H.69 |
| H.31 | Impacts in Washington and Oregon from Shipments of Solid Waste to Hanford | |
| | from Offsite and Shipments of TRU waste to WIPP | H.71 |
| H.32 | Impacts in Washington and Oregon by State from Offsite Shipments of | |
| | Solid Wastes to and from Hanford | H.73 |
| H.33 | Hazardous Chemical Concentrations 100 m (109 yd) Downwind from | |
| | Severe Transportation Accidents | H.75 |
| H.34 | Comparison of Offsite LLW and MLLW Volumes and Impacts Between the WM PEIS | |
| | Information Package and HSW EIS | H.81 |
| H.35 | Comparison of Potential Transportation Impacts for Shipments of TRU Waste from | |
| | Hanford from Hanford to WIPP | H.83 |
| I.1 | Hanford Site Biological Resources Management Resource Levels and Associated | |
| | Definitions | I.3 |
| I.2 | Area C Plant Communities in Which Purple Mat, Crouching Milkvetch, and/or | |
| | Stalked-Pod Milkvetch Were Observed | I.12 |
| I.3 | General Classes of Mitigation Actions and Biological Resource Levels of Concern to | |
| | Which They Apply | I.20 |
| I.4 | Habitats and Species Subject to Mitigation Based on Existing Conditions and | |
| | Current Mitigation Guidelines | I.26 |
| I.5 | Summary of Assumptions Used to Derive Abiotic Media Concentrations Used in | |
| | Hanford and Hanford Plus Background Exposure Scenarios | I.30 |
| I.6 | Hanford and Hanford Plus Background Radionuclide Concentrations in Well Water, | |
| | Pore Water, Sediment, Soil, and River Water for Each Time Period and | |
| | Alternative Group | I.32 |
| I.7 | Hanford and Hanford plus Background Total Uranium Concentrations in Well Water, | |
| | Pore Water, Sediment, Soil, and River Water for Each Time Period and | |
| | Alternative Group | I.39 |
| I.8 | Ecological Contaminant Exposure Model Receptors | I.40 |
| . . | | |
| T 1 | Applicable State Noise Limitations Rased on Source and Recentor FDNA Designation | T 1 |

| J.2 | Estimated Distances of Attenuation by Octave Band for a 132-dBA Diesel Locomotive | J.2 |
|-----|--|------|
| | Previously Conducted Cultural Resources Reviews Cultural Resources Identified in Project Area | |
| L.1 | Comparison of Initial Assessment and HSW EIS Inventories | L.10 |

| L.2 | Comparison of Selected Values of Key Parameters Used in Source Term Release | |
|------|---|------|
| | Models for the System Assessment Capability Analysis Described in this Appendix, | |
| | the HSW EIS Analysis Described in Appendix G, and the Solid Waste Burial Ground | |
| | Performance Assessments for 200 West and East Areas Described by | |
| | Wood et al. (1995) and Wood (1996) | L.22 |
| L.3 | Technetium-99 Distribution Coefficients | L.24 |
| L.4 | Iodine Distribution Coefficients | L.24 |
| L.5 | Uranium Distribution Coefficients | L.25 |
| L.6 | Summary of Hydrogeologic Units Used in This Study | L.26 |
| L.7 | Barrier Design Lifetimes and Estimated Recharge Rates | L.27 |
| L.8 | Estimated Recharge Rates for Predominant Soil Types and Sediments with a | |
| | Shrub-Steppe Plant Community | L.29 |
| L.9 | Estimated Recharge Rates for Native Soils and Backfilled Sediments without Vegetation | L.29 |
| L.10 | Recharge Estimates for the Initial Assessment | L.30 |
| N.1 | Hanford Performance Management Plan Acceleration Goals | N.14 |